

SUMMATION THEOREMS INVOLVING APPELL'S HYPERGEOMETRIC FUNCTIONS F_2

M. S. BABOO^{1,*} AND ASHFAQ AHMAD²

¹School of Basic Sciences and Research, Sharda University, Greater Noida, U. P., 201306, India

²Department of Applied Sciences and Humanities, Faculty of Engineering and Technology, Jamia Millia Islamia (A Central University), New Delhi -110025, India

*Corresponding Author: mesub007@gmail.com

Received Jan. 14, 2019

ABSTRACT. The objective of this paper is to find the closed form of summation theorem for Appell's double hypergeometric function second kind with suitable convergence conditions. In this paper, we find summation theorems of $F_2[A; B, C; A, A; x, y]$, where y takes form $\frac{1-x}{1+x}, \frac{x-1}{2x-1}, \frac{8(1-x)}{8+x}, \frac{1-x}{1+8x}, \frac{x-1}{9x-1}, \frac{1-x}{1+3x}, \frac{x-1}{4x-1}, \frac{3(1-x)}{3+x}, \frac{x-1}{3x-1}, \frac{1-x}{1+2x}$ and other rational functions of x .

2010 Mathematics Subject Classification. Primary 33C65, 33C20; Secondary 33C05.

Key words and phrases. Generalized hypergeometric function; Appell functions of Second kind; Generalized Kummer's First, Second and Third summation theorems.

1. INTRODUCTION, DEFINITIONS AND PRELIMINARIES

In the usual notation, let \mathbb{R} and \mathbb{C} denote the sets of real and complex numbers, respectively. For definitions of Pochhammer symbol, generalized hypergeometric function ${}_pF_q$, we refer monumental work of Srivastava and Manocha [28], Rainville [22] and other notation have their usual meanings.

The Appell's function: The Appell's function of second kind is defined as

$$\begin{aligned}
 F_2[A; B, C; D, G; x, y] &= \sum_{r,s=0}^{\infty} \frac{(A)_{r+s}(B)_r(C)_s}{(D)_r(G)_s} \frac{x^r y^s}{r!s!} \\
 &= \sum_{s=0}^{\infty} \frac{(A)_s(C)_s}{(G)_s} \frac{y^s}{s!} {}_2F_1 \left[\begin{matrix} A+s, B; \\ D; \end{matrix} x \right] \\
 &\left(|x| + |y| < 1, D, G \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}
 \tag{1.1}$$

Classical Gauss summation theorem [28, p.30, Equation 1.2(6)]

$${}_2F_1 \left[\begin{matrix} a, b; \\ d; \end{matrix} 1 \right] = \frac{\Gamma(d)\Gamma(d-a-b)}{\Gamma(d-a)\Gamma(d-b)}, \left(\Re(d-a-b) > 0, d \in \mathbb{C} \setminus \mathbb{Z}_0^- \right).
 \tag{1.2}$$

Kummer's first summation theorem [9, p.852, Equation (1.3)]:

$$(1.3) \quad {}_2F_1 \left[\begin{matrix} a, b; \\ a - b + 1; \end{matrix} -1 \right] = \frac{2^{-a} \sqrt{\pi} \Gamma(a - b + 1)}{\Gamma(\frac{1+a}{2})\Gamma(\frac{a}{2} - b + 1)} = \frac{\Gamma(1 + a - b)\Gamma(1 + \frac{a}{2})}{\Gamma(1 + \frac{a}{2} - b)\Gamma(1 + a)} \cdot \left(a - b \in \mathbb{C} \setminus \mathbb{Z}^-; \Re(b) < 1 \right).$$

In the year 2007, some generalization of the Kummer’s first summation theorem were given by Choi-Rathie and Malani [8, pp. 1523–1524, Equations (2.2), (2.3)], recently some more generalization of the Kummer’s first summation theorem were derived by the Qureshi-Baboo [19, p.14, Equations (3.1), (3.2), (3.3) and (3.4)].

Kummer’s second summation theorem [9, p.852, Equation (1.4)]

$$(1.4) \quad {}_2F_1 \left[\begin{matrix} a, b; \frac{1}{2} \\ \frac{1+a+b}{2}; \end{matrix} \right] = \frac{\Gamma(\frac{1}{2})\Gamma(\frac{1+a+b}{2})}{\Gamma(\frac{a+1}{2})\Gamma(\frac{b+1}{2})}; \quad \left(\frac{1 + a + b}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right).$$

A generalization of the Kummer’s second summation theorem is recorded by Prudnikov *et al.* [18, p.(491), Entry (7.3.7.2)]. In the year 2011, a generalization of the Kummer’s second summation theorem was given by Rakha-Rathie [23, p.827, Theorems (1)], recently some more generalization of the Kummer’s second summation theorem were derived by the Qureshi-Baboo [20, p.48, Equations (3.1) and (3.3)].

Kummer’s third summation theorem [9, p.852, Equation (1.5)]

$$(1.5) \quad {}_2F_1 \left[\begin{matrix} a, 1 - a; \frac{1}{2} \\ c; \end{matrix} \right] = \frac{2^{1-c} \sqrt{\pi} \Gamma(c)}{\Gamma(\frac{c+a}{2})\Gamma(\frac{c+1-a}{2})} = \frac{\Gamma(\frac{c}{2})\Gamma(\frac{c+1}{2})}{\Gamma(\frac{c+a}{2})\Gamma(\frac{c+1-a}{2})}; \quad (c \in \mathbb{C} \setminus \mathbb{Z}_0^-).$$

In the year 2011, some generalization of the Kummer’s third summation theorem were given by Rakha-Rathie [23, p.828, Theorems (6, 5)], recently some more generalization of the Kummer’s third summation theorem were derived by the Qureshi-Baboo [21, Equations (3.3) and (3.5)].

Along with these summation theorems, there are so many summation theorems for Gauss hypergeometric function ${}_2F_1$ with different argument. Some summation theorems are recoded in the monographs of Abramowitz [1], Andrews *et al.* [2], Brychkov [5], Erdélyi *et al.* [10], results conjectured by Gosper and given by Geesel and Stanton [11], results derived by Heymann [12] and [13], Per W. karlsson [14], Kummer [15], Lavoie and Trottier [16], Luke [17], Prudnikov *et al.* [18] and Spiegel [26].

2. SUMMATION THEOREMS DERIVED BY THE REDUCTION FORMULA 2.1

Any values of parameters and variables leading to the result which do not make sense, are tacitly excluded, then we can write the Appell’s function of second kind in the following form

$$(2.1) \quad F_2[A; B, C; A, A; x, y] = (1 - x)^{-B}(1 - y)^{-C} {}_2F_1 \left[\begin{matrix} B, C; \\ A; \end{matrix} \frac{xy}{(1 - x)(1 - y)} \right] \left(|x| < 1, |y| < 1, \left| \frac{xy}{(1 - x)(1 - y)} \right| < 1; A \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = a, B = b, C = c, x = \frac{1}{2}$ and $y = \frac{1}{2}$, use the classical Gauss summation theorem, we get

$$(2.2) \quad F_2 \left[a; b, c; a, a; \frac{1}{2}, \frac{1}{2} \right] = 2^{b+c} \frac{\Gamma(a)\Gamma(a-b-c)}{\Gamma(a-b)\Gamma(a-c)}$$

$$\left(\operatorname{Re}(a-b-c) > 0; a \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{1+b+c}{2}, B = b, C = c$ and $y = \frac{1-x}{1+x}$, use the classical Gauss summation theorem, we get

$$(2.3) \quad F_2 \left[\frac{1+b+c}{2}; b, c; \frac{1+b+c}{2}, \frac{1+b+c}{2}; x, \frac{1-x}{1+x} \right] = \frac{(1+x)^c \Gamma(\frac{1}{2}) \Gamma(\frac{1+b+c}{2})}{(2x)^c (1-x)^b \Gamma(\frac{1+b}{2}) \Gamma(\frac{1+c}{2})}$$

$$\left(0 < x < 1; \frac{1+b+c}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{1+b+c-m}{2}, B = b, C = c$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer's second summation theorem recorded by Prudnikov *et al.* [18, p.491, Entry (7.3.7.2)], we get

$$(2.4) \quad F_2 \left[\frac{1+b+c-m}{2}; b, c; \frac{1+b+c-m}{2}, \frac{1+b+c-m}{2}; x, \frac{1-x}{1+x} \right]$$

$$= \frac{2^{b-1} (1+x)^c \Gamma(\frac{b+c+1-m}{2})}{(1-x)^b (2x)^c \Gamma(b)} \sum_{r=0}^m \left\{ \binom{m}{r} \frac{\Gamma(\frac{b+r}{2})}{\Gamma(\frac{c+1+r-m}{2})} \right\},$$

$$\left(0 < x < 1; b, \frac{1+c+b-m}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).$$

In the equation (2.1) put $A = \frac{1+b+c+m}{2}, B = b, C = c$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer's second summation theorem given by Rakha-Rathie [23, p.827, Theorems (1)], we get

$$(2.5) \quad F_2 \left[\frac{1+b+c+m}{2}; b, c; \frac{1+b+c+m}{2}, \frac{1+b+c+m}{2}; x, \frac{1-x}{1+x} \right]$$

$$= \frac{2^{b-1} (1+x)^c \Gamma(\frac{b+c+1+m}{2}) \Gamma(\frac{c-b+1-m}{2})}{(1-x)^b (2x)^c \Gamma(b) \Gamma(\frac{c-b+1+m}{2})} \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(-1)^r \Gamma(\frac{b+r}{2})}{\Gamma(\frac{c+1+r-m}{2})} \right\},$$

$$\left(0 < x < 1; b, \frac{c+b+1+m}{2}, \frac{c-b+1-m}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).$$

In the equation (2.1) put $A = \frac{b+c-m}{2}, B = b, C = c$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer's second summation theorem given by Qureshi-Baboo [20, p.48, Equation (3.1)], we get

$$(2.6) \quad F_2 \left[\frac{b+c-m}{2}; b, c; \frac{b+c-m}{2}, \frac{b+c-m}{2}; x, \frac{1-x}{1+x} \right]$$

$$= \frac{2^{c-1} (1+x)^c \Gamma(\frac{c+b-m}{2})}{(1-x)^b (2x)^c \Gamma(c)} \sum_{r=0}^m \left\{ \binom{m}{r} \left[\frac{\Gamma(\frac{r+c}{2})}{\Gamma(\frac{b+r-m}{2})} + \frac{\Gamma(\frac{r+c+1}{2})}{\Gamma(\frac{b+r-m+1}{2})} \right] \right\},$$

$$\left(0 < x < 1; c, \frac{c+b-m}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right),$$

In the equation (2.1) put $A = \frac{b+c+m}{2}$, $B = b$, $C = c$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer's second summation theorem given by Qureshi-Baboo [20, p.48, Equation (3.3)], we get

$$(2.7) \quad F_2 \left[\frac{b+c+m}{2}; b, c; \frac{b+c+m}{2}, \frac{b+c+m}{2}; x, \frac{1-x}{1+x} \right] \\ = \frac{2^{b-1}(1+x)^c \Gamma(\frac{c+b-m}{2}) \Gamma(\frac{c-b-m}{2})}{(1-x)^b (2x)^c \Gamma(b) \Gamma(\frac{c-b+m}{2})} \sum_{r=0}^m \left\{ \binom{m}{r} \left[\frac{(-1)^r \Gamma(\frac{r+b}{2})}{\Gamma(\frac{c+r-m}{2})} + \frac{(-1)^r \Gamma(\frac{r+b+1}{2})}{\Gamma(\frac{c+r-m+1}{2})} \right] \right\}, \\ \left(0 < x < 1; b, \frac{c+b+m}{2}, \frac{c-b-m}{2}, \frac{c-b+m}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right),$$

In the equation (2.1) put $A = a$, $B = b$, $C = 1 - b$ and $y = \frac{1-x}{1+x}$, use the Kummer's third summation theorem, we get

$$(2.8) \quad F_2 \left[a; b, 1 - b; a, a; x, \frac{1-x}{1+x} \right] = \frac{(1+x)^{1-b} \Gamma(\frac{a}{2}) \Gamma(\frac{a+1}{2})}{(1-x)^b (2x)^{1-b} \Gamma(\frac{a+b}{2}) \Gamma(\frac{a-b+1}{2})} \\ \left(0 < x < 1; a \in \mathbb{C} \setminus \mathbb{Z}_0^- \right),$$

In the equation (2.1) put $A = a$, $B = b$, $C = 1 - b - m$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer's third summation theorem given by Rakha-Rathie [23, p.828, Theorem (6)], we get

$$(2.9) \quad F_2 \left[a; b, 1 - b - m; a, a; x, \frac{1-x}{1+x} \right] \\ = \frac{2^{1-m-a} (1+x)^{1-b-m} \Gamma(\frac{1}{2}) \Gamma(a)}{(1-x)^b (2x)^{1-b-m} \Gamma(\frac{a-b}{2}) \Gamma(\frac{a-b+1}{2})} \sum_{r=0}^m \left\{ \binom{m}{r} \frac{\Gamma(\frac{a-b+r}{2})}{\Gamma(\frac{a+b+r}{2})} \right\} \\ \left(0 < x < 1; a, a - b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right);$$

In the equation (2.1) put $A = a$, $B = b$, $C = 1 - b + m$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer's third summation theorem given by Rakha-Rathie [23, p.828, Theorem (5)], we get

$$(2.10) \quad F_2 \left[a; b, 1 - b + m; a, a; x, \frac{1-x}{1+x} \right] \\ = \frac{2^{1+m-a} (1+x)^{1-b+m} \Gamma(\frac{1}{2}) \Gamma(a) \Gamma(b-m)}{(2x)^{1-b+m} (1-x)^b \Gamma(b) \Gamma(\frac{a-b}{2}) \Gamma(\frac{a-b+1}{2})} \sum_{r=0}^m \left\{ (-1)^r \binom{m}{r} \frac{\Gamma(\frac{a-b+r}{2})}{\Gamma(\frac{a+b+r}{2} - m)} \right\} \\ \left(0 < x < 1; a, b, b - m, a - b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right);$$

In the equation (2.1) put $A = a$, $B = b$, $C = -b - m$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer's third summation theorem given by Qureshi-Baboo [21, p.144, Equation (3.3)], we get

$$F_2 \left[a; b, -b - m; a, a; x, \frac{1-x}{1+x} \right]$$

$$(2.11) \quad = \frac{2^{-1-m-b}(2x)^{b+m}\Gamma(a)}{(1-x)^b(1+x)^{b+m}\Gamma(a-b)} \sum_{r=0}^m \left\{ \binom{m}{r} \left[\frac{\Gamma(\frac{a-b+r}{2})}{\Gamma(\frac{a+b+r}{2})} + \frac{\Gamma(\frac{a-b+r+1}{2})}{\Gamma(\frac{a+b+r+1}{2})} \right] \right\} \\ \left(0 < x < 1; a, a-b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right);$$

In the equation (2.1) put $A = a, B = b, C = -b + m$ and $y = \frac{1-x}{1+x}$, using the generalization of Kummer’s third summation theorem given by Qureshi-Baboo [21, p.145, Equation (3.5)], we get

$$(2.12) \quad F_2 \left[a; b, -b + m; a, a; x, \frac{1-x}{1+x} \right] \\ = \frac{(x)^{b-m}\Gamma(a)\Gamma(b-m)}{2(1-x)^b(1+x)^{b-m}\Gamma(b)\Gamma(a-b)} \sum_{r=0}^m \left\{ \binom{m}{r} (-1)^r \left[\frac{\Gamma(\frac{a-b+r}{2})}{\Gamma(\frac{a+b+r-2m}{2})} + \frac{\Gamma(\frac{a-b+r+1}{2})}{\Gamma(\frac{a+b+r-2m+1}{2})} \right] \right\} \\ \left(0 < x < 1; a, a-b, b-m \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right);$$

In the equation (2.1) put $A = 1 + b - c, B = b, C = c$ and $y = \frac{x-1}{2x-1}$, use the Kummer’s first summation theorem, we get

$$(2.13) \quad F_2 \left[1 + b - c; b, c; 1 + b - c, 1 + b - c; x, \frac{x-1}{2x-1} \right] = \frac{(2x-1)^c \Gamma(1+b-c)\Gamma(1+\frac{b}{2})}{x^c(1-x)^b \Gamma(1+b)\Gamma(1+\frac{b}{2}-c)} \\ \left(\Re(c) < 1; \frac{2}{3} < x < 1; 1+b-c \in \mathbb{C} \setminus \mathbb{Z}_0^- \right);$$

In the equation (2.1) put $A = 1 + b - c - m, B = b, C = c$ and $y = \frac{x-1}{2x-1}$, using the generalization of Kummer’s first summation theorem given by Choi, Rathie and Malani [8, p.1524, Equation (2.2)], we get

$$(2.14) \quad F_2 \left[1 + b - c - m; b, c; 1 + b - c - m, 1 + b - c - m; x, \frac{x-1}{2x-1} \right] \\ = \frac{(2x-1)^c \Gamma(1+b-c-m)}{2(x)^c(1-x)^b \Gamma(b)} \sum_{r=0}^m \left\{ \binom{m}{r} \frac{\Gamma(\frac{r+b}{2})}{\Gamma(\frac{r+b}{2} + 1 - c - m)} \right\} \\ \left(\Re(c) < (\frac{2-m}{2}); \frac{2}{3} < x < 1; b, 1+b-c-m \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right),$$

In the equation (2.1) put $A = 1 + b - c + m, B = b, C = c$ and $y = \frac{x-1}{2x-1}$, using the generalization of Kummer’s first summation theorem given by Choi, Rathie and Malani [8, p.1524, Equation (2.3)], we get

$$(2.15) \quad F_2 \left[1 + b - c + m; b, c; 1 + b - c + m, 1 + b - c + m; x, \frac{x-1}{2x-1} \right] \\ = \frac{(2x-1)^c \Gamma(1+b-c+m)}{2(x)^c(1-x)^b \Gamma(b)(1-c)_m} \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(-1)^r \Gamma(\frac{r+b}{2})}{\Gamma(\frac{r+b}{2} + 1 - c)} \right\} \\ \left(\Re(c) < (\frac{m+2}{2}); \frac{2}{3} < x < 1; b, 1-c, 1+b-c+m \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).$$

In the equation (2.1) put $A = b - c - m, B = b, C = c$ and $y = \frac{x-1}{2x-1}$, use the summation theorem given by Qureshi-Baboo [19, p.14, Equation (3.1)], we get

$$\begin{aligned}
 &F_2 \left[b - c - m; b, c; b - c - m, b - c - m; x, \frac{x - 1}{2x - 1} \right] \\
 (2.16) \quad &= \frac{(2x - 1)^c \Gamma(b - c - m)}{2 (x)^c (1 - x)^b \Gamma(b)} \sum_{r=0}^m \left\{ \binom{m}{r} \left[\frac{\Gamma(\frac{b+r}{2})}{\Gamma(\frac{b+r-2c-2m}{2})} + \frac{\Gamma(\frac{b+r+1}{2})}{\Gamma(\frac{b+r+1-2c-2m}{2})} \right] \right\}, \\
 &\left(\Re(c) < \left(\frac{1-m}{2}\right); \frac{2}{3} < x < 1; b, b - c - m \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right),
 \end{aligned}$$

In the equation (2.1) put $A = b - c + m, B = b, C = c$ and $y = \frac{x-1}{2x-1}$, use the summation theorem given by Qureshi-Baboo [19, p.14, Equation (3.2)], we get

$$\begin{aligned}
 &F_2 \left[b - c + m; b, c; b - c + m, b - c + m; x, \frac{x - 1}{2x - 1} \right] \\
 (2.17) \quad &= \frac{(2x - 1)^c \Gamma(b - c + m)}{2 (x)^c (1 - x)^b \Gamma(b) (-c)_m} \sum_{r=0}^m \left\{ \binom{m}{r} \left[\frac{(-1)^r \Gamma(\frac{b+r}{2})}{\Gamma(\frac{b+r-2c}{2})} + \frac{(-1)^r \Gamma(\frac{b+r+1}{2})}{\Gamma(\frac{b+r+1-2c}{2})} \right] \right\}, \\
 &\left(\Re(c) < \left(\frac{m+1}{2}\right); \frac{2}{3} < x < 1; b, -c, b - c + m \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).
 \end{aligned}$$

In the equation (2.1) put $A = \frac{a+4-\sqrt{2-a}}{2}, B = a, C = \frac{a-2-\sqrt{2-a}}{2}$ and $y = \frac{x-1}{2x-1}$ and using a result of Brychkov [5, p.579, Equation (100)], we get

$$\begin{aligned}
 &F_2 \left[\frac{a + 4 - \sqrt{2 - a}}{2}; a, \frac{a - 2 - \sqrt{2 - a}}{2}; \frac{a + 4 - \sqrt{2 - a}}{2}, \frac{a + 4 - \sqrt{2 - a}}{2}; x, \frac{x - 1}{2x - 1} \right] = \\
 (2.18) \quad &= \left(\frac{x}{2x - 1} \right)^{\frac{2-a+\sqrt{2-a}}{2}} \frac{2 + a(3 + \sqrt{2 - a})}{(1 - x)^a 2^{a+1}} \\
 &\left(\frac{2}{3} < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}$$

In the equation (2.1) put $A = \frac{a+5-\sqrt{7-3a}}{2}, B = a, C = \frac{a-3-\sqrt{7-3a}}{2}$ and $y = \frac{x-1}{2x-1}$ and using a result of Brychkov [5, p.579, Equation (101)], we get

$$\begin{aligned}
 &F_2 \left[\frac{a + 5 - \sqrt{7 - 3a}}{2}; a, \frac{a - 3 - \sqrt{7 - 3a}}{2}; \frac{a + 5 - \sqrt{7 - 3a}}{2}, \frac{a + 5 - \sqrt{7 - 3a}}{2}; x, \frac{x - 1}{2x - 1} \right] = \\
 (2.19) \quad &= \left(\frac{x}{2x - 1} \right)^{\frac{3-a+\sqrt{7-3a}}{2}} \frac{6 + a(15 - a + 4\sqrt{7 - 3a})}{6(1 - x)^a 2^{a+1}} \\
 &\left(\frac{2}{3} < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}$$

In the equation (2.1) put $A = \frac{2a+2}{3}, B = \frac{a}{2}, C = \frac{a+1}{2}$ and $y = \frac{8(1-x)}{x+8}$, using a result of Andrews *et al.* [2, p.131, Entry 3.1.20], Kummer [15, p.136, Article 25(6)] see also Prudnikov *et al.* [18, p.495, Equation (38)], we get

$$(2.20) \quad F_2 \left[\frac{2a + 2}{3}; \frac{a}{2}, \frac{a + 1}{2}; \frac{2a + 2}{3}, \frac{2a + 2}{3}; x, \frac{8(1 - x)}{x + 8} \right] = \left(\frac{3}{2} \right)^a \frac{(x + 8)^{\frac{a+1}{2}} \sqrt{\pi} \Gamma(\frac{2a+2}{3})}{(1 - x)^{\frac{a}{2}} (9x)^{\frac{a+1}{2}} \Gamma(\frac{a+4}{6}) \Gamma(\frac{a+1}{2})}$$

$$\left(0 < x < 1; \frac{2a+2}{3} \in \mathbb{C} \setminus \mathbb{Z}_0^-\right)$$

In the equation (2.1) put $A = \frac{2}{3}, B = -a, C = 2a + 1$ and $y = \frac{8(1-x)}{x+8}$ and using a result of Heymann, see Per W. Karlsson [14, p.335, Equation (3.2)], see also Brychkov [5, p.584, Equation (144)], we get

$$(2.21) \quad F_2 \left[\frac{2}{3}; -a, 2a + 1; \frac{2}{3}, \frac{2}{3}; x, \frac{8(1-x)}{x+8} \right] = \frac{2 \times 3^a (1-x)^a (x+8)^{2a+1} \sin(\pi a + \frac{5\pi}{6})}{(9x)^{2a+1}}$$

$$\left(0 < x < 1\right)$$

In the equation (2.1) put $A = \frac{4}{3}, B = -a, C = 2a + 2$ and $y = \frac{8(1-x)}{x+8}$ and using a result of Heymann, see Per W. Karlsson [14, p.335, Equation (3.3)], see also Brychkov [5, p.584, Equation (145)], we get

$$(2.22) \quad F_2 \left[\frac{4}{3}; -a, 2a + 2; \frac{4}{3}, \frac{4}{3}; x, \frac{8(1-x)}{x+8} \right] = \frac{3^a (1-x)^a (x+8)^{2a+2} \Gamma(\frac{3}{2}) \Gamma(\frac{1}{6})}{(9x)^{2a+2} \Gamma(\frac{1}{6} - a) \Gamma(a + \frac{3}{2})}$$

$$\left(0 < x < 1\right)$$

In the equation (2.1) put $A = 4a + \frac{1}{3}, B = 3a, C = 3a + \frac{1}{4}$ and $y = \frac{8(1-x)}{x+8}$ and using a summation formula conjectured by Gosper, see Per W. Karlsson [14, p.335, Equation (3.4)], see also Brychkov [5, p.584, Equation (146)], we get

$$(2.23) \quad F_2 \left[4a + \frac{1}{3}; 3a, 3a + \frac{1}{4}; 4a + \frac{1}{3}, 4a + \frac{1}{3}; x, \frac{8(1-x)}{x+8} \right] =$$

$$= \frac{(108)^a (x+8)^{3a+\frac{1}{4}} \Gamma(a + \frac{7}{12}) \Gamma(a + \frac{5}{6}) \Gamma(\frac{3}{4}) \Gamma(\frac{2}{3})}{(1-x)^{3a} (9x)^{3a+\frac{1}{4}} \Gamma(a + \frac{3}{4}) \Gamma(a + \frac{2}{3}) \Gamma(\frac{7}{12}) \Gamma(\frac{5}{6})}$$

$$\left(0 < x < 1; 4a + \frac{1}{3}, a + \frac{7}{12}, a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^-\right)$$

In the equation (2.1) put $A = 4a + \frac{1}{3}, B = 3a, C = 3a - \frac{1}{4}$ and $y = \frac{8(1-x)}{x+8}$ and using a summation formula conjectured by Gosper, see Per W. Karlsson [14, p.335, Equation (3.5)], see also Brychkov [5, p.584, Equation (147)], we get

$$(2.24) \quad F_2 \left[4a + \frac{1}{3}; 3a, 3a - \frac{1}{4}; 4a + \frac{1}{3}, 4a + \frac{1}{3}; x, \frac{8(1-x)}{x+8} \right] =$$

$$= \frac{(108)^a (x+8)^{3a-\frac{1}{4}} \Gamma(a + \frac{1}{12}) \Gamma(a + \frac{5}{6}) \Gamma(\frac{1}{4}) \Gamma(\frac{2}{3})}{(1-x)^{3a} (9x)^{3a-\frac{1}{4}} \Gamma(a + \frac{1}{4}) \Gamma(a + \frac{2}{3}) \Gamma(\frac{1}{12}) \Gamma(\frac{5}{6})}$$

$$\left(0 < x < 1; 4a + \frac{1}{3}, a + \frac{1}{12}, a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^-\right)$$

In the equation (2.1) put $A = 4a + \frac{2}{3}, B = 3a, C = 3a + \frac{1}{2}$ and $y = \frac{8(1-x)}{x+8}$ and using a result of Kummer [15, p.136, Article 25(6)], see also Per W. Karlsson [14, p.335, Equation (3.1)], see also Brychkov [5, p.584, Equation (143)], we get

$$\begin{aligned}
 (2.25) \quad & F_2 \left[4a + \frac{2}{3}; 3a, 3a + \frac{1}{2}; 4a + \frac{2}{3}, 4a + \frac{2}{3}; x, \frac{8(1-x)}{x+8} \right] = \\
 & = \frac{(27)^a (9x)^{3a+\frac{1}{2}} \Gamma(2a + \frac{5}{6}) \Gamma(\frac{1}{2})}{(x+8)^{3a+\frac{1}{2}} (1-x)^{3a} \Gamma(a + \frac{5}{6}) \Gamma(a + \frac{1}{2})} \\
 & \quad \left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}$$

In the equation (2.1) put $A = 4a, B = a, C = a + \frac{1}{2}$ and $y = \frac{8(1-x)}{x+8}$ and using a result of Prudnikov *et al.* [18, p.496, Equation (41)], we get

$$\begin{aligned}
 (2.26) \quad & F_2 \left[4a; a, a + \frac{1}{2}; 4a, 4a; x, \frac{8(1-x)}{x+8} \right] = \frac{(3)^{2a} (x+8)^{a+\frac{1}{2}} \Gamma(\frac{1}{2}) \Gamma(4a)}{(9x)^{a+\frac{1}{2}} (1-x)^a \Gamma(3a) \Gamma(a + \frac{1}{2}) 2^{6a-1}} \\
 & \quad \left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}$$

In the equation (2.1) put $A = \frac{2a+5}{6}, B = \frac{a}{2}, C = \frac{a+1}{2}$ and $y = \frac{1-x}{1+8x}$, using a result of Andrews *et al.* [2, p.131, Entry 3.1.17]; Abramowitz [1, p.557, Entry 15.1.30]; Kummer [15, p.135, Article 25(4)]; Per W. Karlsson [14, p.330, Equation (1.1)] and Brychkov [5, p.580, Equation (115)], we get

$$\begin{aligned}
 (2.27) \quad & F_2 \left[\frac{2a+5}{6}; \frac{a}{2}, \frac{a+1}{2}; \frac{2a+5}{6}, \frac{2a+5}{6}; x, \frac{1-x}{1+8x} \right] = \left(\frac{3}{4} \right)^a \frac{\sqrt{\pi} (1+8x)^{\frac{a+1}{2}} \Gamma(\frac{2a+2}{3})}{(1-x)^{\frac{a}{2}} (9x)^{\frac{a+1}{2}} \Gamma(\frac{a+4}{6}) \Gamma(\frac{a+1}{2})} \\
 & \quad \left(0 < x < 1; \frac{2a+5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}$$

In the equation (2.1) put $A = a + \frac{5}{6}, B = \frac{1-2a}{2}, C = 2a$ and $y = \frac{1-x}{1+8x}$ and using a result of Heymann, see Per.W. Karlsson [14, p.330, Equation (1.2)] and Brychkov [5, p.580, Equation 116], we get

$$\begin{aligned}
 (2.28) \quad & F_2 \left[a + \frac{5}{6}; \frac{1-2a}{2}, 2a; a + \frac{5}{6}, a + \frac{5}{6}; x, \frac{1-x}{1+8x} \right] = \frac{3^a (1+8x)^{2a} (1-x)^{\frac{2a-1}{2}} \Gamma(a + \frac{5}{6}) \Gamma(\frac{2}{3})}{4^a (9x)^{2a} \Gamma(a + \frac{2}{3}) \Gamma(\frac{5}{6})} \\
 & \quad \left(0 < x < 1; a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}$$

In the equation (2.1) put $A = a + \frac{2}{3}, B = 1 - a, C = 2a$ and $y = \frac{1-x}{1+8x}$ and using a result of Heymann, see Per W. Karlsson [14, p.330, Equation (1.3)], see also Brychkov [5, p.581, Equation (117)], we get

$$\begin{aligned}
 (2.29) \quad & F_2 \left[a + \frac{2}{3}; 1 - a, 2a; a + \frac{2}{3}, a + \frac{2}{3}; x, \frac{1-x}{1+8x} \right] = \frac{3^a (1-x)^{a-1} (1+8x)^{2a} \Gamma(a + \frac{2}{3}) \Gamma(\frac{1}{2})}{4^a (9x)^{2a} \Gamma(a + \frac{1}{2}) \Gamma(\frac{2}{3})} \\
 & \quad \left(0 < x < 1; a + \frac{2}{3} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)
 \end{aligned}$$

In the equation (2.1) put $A = 2a + \frac{5}{4}, B = \frac{1}{4} - a, C = -a$ and $y = \frac{1-x}{1+8x}$ and using a summation formula conjectured by Gosper, see Per W. Karlsson [14, p.330, Equation (1.4)], see also Brychkov [5, p.581, Equation (118)], we get

$$(2.30) \quad F_2 \left[2a + \frac{5}{4}; \frac{1}{4} - a, -a; 2a + \frac{5}{4}, 2a + \frac{5}{4}; x, \frac{1-x}{1+8x} \right] = \frac{2^{6a}(1-x)^{a-\frac{1}{4}}(9x)^a \Gamma(2a + \frac{5}{4}) \Gamma(\frac{2}{3}) \Gamma(\frac{13}{12})}{3^{5a}(1+8x)^a \Gamma(a + \frac{2}{3}) \Gamma(a + \frac{13}{12}) \Gamma(\frac{5}{4})}$$

$$\left(0 < x < 1; 2a + \frac{5}{4} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = 2a + \frac{5}{4}, B = \frac{1}{4} - a, C = -a$ and $y = \frac{1-x}{1+8x}$ and using a summation formula conjectured by Gosper, see Per W. Karlsson [14, p.330, Equation (1.5)], see also Brychkov [5, p.581, Equation (119)], we get

$$(2.31) \quad F_2 \left[2a + \frac{9}{4}; \frac{1}{4} - a, -a; 2a + \frac{9}{4}, 2a + \frac{9}{4}; x, \frac{1-x}{1+8x} \right] = \frac{2^{6a}(1-x)^{a-\frac{1}{4}}(9x)^a \Gamma(2a + \frac{9}{4}) \Gamma(\frac{4}{3}) \Gamma(\frac{17}{12})}{3^{5a}(1+8x)^a \Gamma(a + \frac{4}{3}) \Gamma(a + \frac{17}{12}) \Gamma(\frac{9}{4})}$$

$$\left(0 < x < 1; 2a + \frac{9}{4} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{4}{3} - a, B = a, C = 1 - 2a$ and $y = \frac{1-x}{8x+1}$ and using a result of Prudnikov *et al.* [18, p.495, Equation (27)], we get

$$(2.32) \quad F_2 \left[\frac{4}{3} - a; a, 1 - 2a; \frac{4}{3} - a, \frac{4}{3} - a; x, \frac{1-x}{8x+1} \right] = \left(\frac{9x}{8x+1} \right)^{2a-1} \frac{\Gamma(\frac{2}{3} - a) \Gamma(\frac{4}{3} - a)}{(1-x)^a 3^a \Gamma(\frac{2}{3}) \Gamma(\frac{4}{3} - 2a)}$$

$$\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{2a+5}{6}, B = \frac{a}{2}, C = \frac{2-a}{6}$ and $y = \frac{x-1}{9x-1}$, using a result of Andrews *et al.* [2, p.177, Question 3(a)], see also Prudnikov *et al.* [18, p.494, Equation (19)], we get

$$(2.33) \quad F_2 \left[\frac{2a+5}{6}; \frac{a}{2}, \frac{2-a}{6}; \frac{2a+5}{6}, \frac{2a+5}{6}; x, \frac{x-1}{9x-1} \right] = \frac{\sqrt{\pi}(9x-1)^{\frac{2-a}{6}} \Gamma(\frac{2a+2}{3})}{2^{\frac{a}{2}}(1-x)^{\frac{a}{2}}(8x)^{\frac{2-a}{6}} \Gamma(\frac{a+4}{6}) \Gamma(\frac{a+1}{2})}$$

$$\left(0 < x < 1; \frac{2a+5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = -a + \frac{4}{3}, B = a, C = a + \frac{1}{3}$ and $y = \frac{x-1}{9x-1}$, using a result of Lavoie and Trottier [16, p.45, Equation (8)], see also Prudnikov *et al.* [18, p.494, Equation (18)], we get

$$(2.34) \quad F_2 \left[-a + \frac{4}{3}; a, a + \frac{1}{3}; -a + \frac{4}{3}, -a + \frac{4}{3}; x, \frac{x-1}{9x-1} \right] = \left(\frac{2}{3} \right)^{3a} \frac{(9x-1)^{a+\frac{1}{3}} \Gamma(\frac{2}{3} - a) \Gamma(\frac{4}{3} - a)}{(8x)^{a+\frac{1}{3}}(1-x)^a \Gamma(\frac{2}{3}) \Gamma(\frac{4}{3} - 2a)}$$

$$\left(0 < x < 1; -a + \frac{4}{3}, -a + \frac{2}{3} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = 2b + \frac{1}{2}, B = \frac{1}{2}, C = 1 - b$ and $y = \frac{1-x}{3x+1}$, using a result of Spiegel [26, p.894]; Luke [17, p.273, Equation 6.8(20)], we get

$$(2.35) \quad F_2 \left[2b + \frac{1}{2}; \frac{1}{2}, 1 - b; 2b + \frac{1}{2}, 2b + \frac{1}{2}; x, \frac{1-x}{3x+1} \right] = \frac{2(4x)^{b-1} \Gamma(b) \Gamma(2b + \frac{1}{2})}{3(1-x)^{\frac{1}{2}}(3x-1)^{b-1} \Gamma(2b) \Gamma(b + \frac{1}{2})}$$

$$\left(0 < x < 1; b, 2b + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{5}{2} - 2a, B = \frac{1}{2}, C = a$ and $y = \frac{1-x}{3x+1}$ and using a result of Prudnikov *et al.* [18, p.495, Equation (31)], we get

$$(2.36) \quad F_2 \left[\frac{5}{2} - 2a; \frac{1}{2}, a; \frac{5}{2} - 2a, \frac{5}{2} - 2a; x, \frac{1-x}{3x+1} \right] = \left(\frac{3x+1}{4x} \right)^a \frac{2^{2a} \Gamma(\frac{1}{2}) \Gamma(\frac{5}{2} - 2a)}{3(1-x)^{\frac{1}{2}} \Gamma(\frac{3}{2} - a) \Gamma(\frac{3}{2} - a)}$$

$$\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = -2a + \frac{3}{2}, B = a, C = a + \frac{1}{2}$ and $y = \frac{x-1}{4x-1}$, using a result of Abramowitz [1, p.557, Entry 15.1.29]; Erdélyi *et al.* [10, p.104, Entry 2.8.53], see also Per W. Karlsson [14, p.334, Equation (2.12)], Prudnikov *et al.* [18, p.494, Equation (15)], we get

$$(2.37) \quad F_2 \left[-2a + \frac{3}{2}; a, a + \frac{1}{2}; -2a + \frac{3}{2}, -2a + \frac{3}{2}; x, \frac{x-1}{4x-1} \right] =$$

$$= \left(\frac{9}{8} \right)^{2a} \frac{(4x-1)^{a+\frac{1}{2}} \Gamma(\frac{4}{3}) \Gamma(-2a + \frac{3}{2})}{(1-x)^a (3x)^{a+\frac{1}{2}} \Gamma(\frac{3}{2}) \Gamma(-2a + \frac{4}{3})}$$

$$\left(0 < x < 1; -2a + \frac{3}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = a + \frac{1}{2}, B = a, C = 1 - \frac{a}{2}$ and $y = \frac{x-1}{4x-1}$ and using a result of Prudnikov *et al.* [18, p.494, Equation (14)], we get

$$(2.38) \quad F_2 \left[a + \frac{1}{2}; a, 1 - \frac{a}{2}; a + \frac{1}{2}, a + \frac{1}{2}; x, \frac{x-1}{4x-1} \right] = \left(\frac{x}{4x-1} \right)^{\frac{a}{2}-1} \frac{\Gamma(\frac{1}{2}) \Gamma(\frac{2a+1}{2})}{3(1-x)^a \Gamma(\frac{a+1}{2}) \Gamma(\frac{a+1}{2})}$$

$$\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = 3a, B = b, C = \frac{1}{2}$ and $y = \frac{3(1-x)}{x+3}$ and using a result of Luke [17, p.273, Equation 6.8(18)], see also Prudnikov *et al.* [18, p.495, Equation (35)], we get

$$(2.39) \quad F_2 \left[3a; b, \frac{1}{2}; 3a, 3a; x, \frac{3(1-x)}{x+3} \right] = \left(\frac{16}{27} \right)^a \frac{(x+3)^{\frac{1}{2}} \Gamma(a) \Gamma(3a)}{(4x)^{\frac{1}{2}} (1-x)^a (\Gamma(2a))^2}$$

$$\left(0 < x < 1; a \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{5-a}{2}, B = 2, C = a$ and $y = \frac{x-1}{3x-1}$ and using a result of Prudnikov *et al.* [18, p.494, Equation (13)], we get

$$(2.40) \quad F_2 \left[\frac{5-a}{2}; 2, a; \frac{5-a}{2}, \frac{5-a}{2}; x, \frac{x-1}{3x-1} \right] = \left(\frac{3x-1}{2x} \right)^a \frac{3-a}{3(1-x)^2}$$

$$\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = a + 2, B = a, C = 1 - 2a$ and $y = \frac{1-x}{1+2x}$ and using a result of Prudnikov *et al.* [18, p.495, Equation (34)], we get

$$(2.41) \quad F_2 \left[a + 2; a, 1 - 2a; a + 2, a + 2; x, \frac{1 - x}{1 + 2x} \right] = \left(\frac{2x}{1 + 2x} \right)^{2a-1} \frac{2(a + 1)}{3(1 - x)^a} \\ \left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = a + \frac{3}{4}, B = 2a, C = a + \frac{1}{4}$ and $y = \left(\frac{(\sqrt{2}-1)(1-x)}{2x+\sqrt{2}-1} \right)$, using a result of Andrews et al. [2, p.177, Question 3(b)], we get

$$(2.42) \quad F_2 \left[a + \frac{3}{4}; 2a, a + \frac{1}{4}; a + \frac{3}{4}, a + \frac{3}{4}; x, \frac{(\sqrt{2} - 1)(1 - x)}{2x + \sqrt{2} - 1} \right] \\ = \frac{\sqrt{\pi} \Gamma(a + \frac{3}{4})(2x + \sqrt{2} - 1)^{a+\frac{1}{4}}}{[x(1 + \sqrt{2})]^{a+\frac{1}{4}}(1 - x)^{2a}(4 - 2\sqrt{2})^{2a} \Gamma(\frac{2a+3}{4}) \Gamma(\frac{a+1}{2})} \\ \left(0 < x < 1; a + \frac{3}{4} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{2a+3}{4}, B = \frac{1}{2}, C = a$ and $y = \frac{(1-\sqrt{2})(1-x)}{2x+(1-\sqrt{2})(1-x)}$ and using a result of Prudnikov et al. [18, p.494, Equation (16)], we get

$$(2.43) \quad F_2 \left[\frac{2a + 3}{4}; \frac{1}{2}, a; \frac{2a + 3}{4}, \frac{2a + 3}{4}; x, \frac{(1 - \sqrt{2})(1 - x)}{2x + (1 - \sqrt{2})(1 - x)} \right] = \\ = \left(\frac{2x + (1 - \sqrt{2})(1 - x)}{2x} \right)^a \frac{\Gamma(\frac{1}{2}) \Gamma(\frac{2a+3}{4})}{2^{\frac{a}{2}} (1 - x)^{\frac{1}{2}} \Gamma(\frac{a+2}{4}) \Gamma(\frac{a+3}{4})} \\ \left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{2a+5}{6}, B = a, C = \frac{2-a}{3}$ and $y = \frac{(4-3\sqrt{2})(1-x)}{8x+(4-3\sqrt{2})(1-x)}$ and using a result of Prudnikov et al. [18, p.495, Equation (22)], we get

$$(2.44) \quad F_2 \left[\frac{2a + 5}{6}; a, \frac{2 - a}{3}; \frac{2a + 5}{6}, \frac{2a + 5}{6}; x, \frac{(4 - 3\sqrt{2})(1 - x)}{8x + (4 - 3\sqrt{2})(1 - x)} \right] = \\ = \left(\frac{8x}{8x + (4 - 3\sqrt{2})(1 - x)} \right)^{\frac{a-2}{3}} \left(\frac{2}{3} \right)^{\frac{a}{2}} \frac{\Gamma(\frac{1}{2}) \Gamma(\frac{2a+5}{6})}{(1 - x)^a \Gamma(\frac{a+3}{6}) \Gamma(\frac{a+5}{6})} \\ \left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{3-2a}{2}, B = a, C = 2 - 3a$ and $y = \frac{(2-\sqrt{3})(1-x)}{4x+(2-\sqrt{3})(1-x)}$ and using a result of Prudnikov et al. [18, p.495, Equation (25)], we get

$$(2.45) \quad F_2 \left[\frac{3 - 2a}{2}; a, 2 - 3a; \frac{3 - 2a}{2}, \frac{3 - 2a}{2}; x, \frac{(2 - \sqrt{3})(1 - x)}{4x + (2 - \sqrt{3})(1 - x)} \right] = \\ = \left(\frac{4x}{4x + (2 - \sqrt{3})(1 - x)} \right)^{3a-2} \frac{3^{\frac{3a}{2}} \Gamma(\frac{4}{3}) \Gamma(\frac{3-2a}{2})}{(1 - x)^a 2^{2a-1} \Gamma(\frac{1}{2}) \Gamma(\frac{4-3a}{3})}$$

$$\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right)$$

In the equation (2.1) put $A = \frac{2a+1}{2}, B = a, C = \frac{2a+1}{4}$ and $y = \frac{(2\sqrt{2}-2)(1-x)}{x+(2\sqrt{2}-2)(1-x)}$ and using a result of Prudnikov *et al.* [18, p.495, Equation (36)], we get

$$\begin{aligned} &F_2 \left[\frac{2a+1}{2}; a, \frac{2a+1}{4}; \frac{2a+1}{2}, \frac{2a+1}{2}; x, \frac{(2\sqrt{2}-2)(1-x)}{x+(2\sqrt{2}-2)(1-x)} \right] = \\ (2.46) \quad &= \left(\frac{x+(2\sqrt{2}-2)(1-x)}{x} \right)^{\frac{2a+1}{4}} \left(\frac{2+\sqrt{2}}{2} \right)^a \frac{\Gamma(\frac{1}{2})\Gamma(\frac{3+2a}{4})}{(1-x)^a \Gamma(\frac{a+2}{4})\Gamma(\frac{a+3}{4})} \\ &\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right) \end{aligned}$$

In the equation (2.1) put $A = \frac{4a+1}{3}, B = a, C = \frac{4a+1}{6}$ and $y = \frac{(12\sqrt{2}-16)(1-x)}{x+(12\sqrt{2}-16)(1-x)}$ and using a result of Prudnikov *et al.* [18, p.496, Equation (43)], see also Brychkov [5, p.587, Equation (172)], we get

$$\begin{aligned} &F_2 \left[\frac{4a+1}{3}; a, \frac{4a+1}{6}; \frac{4a+1}{3}, \frac{4a+1}{3}; x, \frac{(12\sqrt{2}-16)(1-x)}{x+(12\sqrt{2}-16)(1-x)} \right] = \\ (2.47) \quad &= \left(\frac{x+(12\sqrt{2}-16)(1-x)}{x} \right)^{\frac{4a+1}{6}} \left(\frac{2+\sqrt{2}}{2} \right)^{2a} \frac{\Gamma(\frac{1}{2})\Gamma(\frac{2a+2}{3})}{(1-x)^a \Gamma(\frac{a+1}{2})\Gamma(\frac{a+4}{6})} \\ &\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right) \end{aligned}$$

In the equation (2.1) put $A = 4a - 1, B = a, C = \frac{4a-1}{2}$ and $y = \frac{(12\sqrt{2}-16)(1-x)}{x+(12\sqrt{2}-16)(1-x)}$ and using a result of Prudnikov *et al.* [18, p.496, Equation (46)], we get

$$\begin{aligned} &F_2 \left[4a - 1; a, \frac{4a-1}{2}; 4a - 1, 4a - 1; x, \frac{(12\sqrt{2}-16)(1-x)}{x+(12\sqrt{2}-16)(1-x)} \right] = \\ (2.48) \quad &= \left(\frac{x+(12\sqrt{2}-16)(1-x)}{x} \right)^{\frac{4a-1}{2}} \left(\frac{2+\sqrt{2}}{2} \right)^{2a} \frac{\Gamma(a)}{(1-x)^a \Gamma(\frac{3a}{2})} \\ &\left(0 < x < 1; 4a + \frac{2}{3}, 2a + \frac{5}{6} \in \mathbb{C} \setminus \mathbb{Z}_0^- \right) \end{aligned}$$

REFERENCES

[1] Abramowitz, M. and Stegun, I. A. ; *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards Applied Mathematics Series, vol. 55, U. S. Government Printing Office, Washington, 1964.

[2] Andrews, G. E., Askey, R. and Roy, R. ; *Special Function, Encyclopedia of Mathematics and its Applications*, Vol. 71, Cambridge University Press, Cambridge, 1999.

[3] Appell, P. and Kampe de Fariet, J. ; *Fonctions Hypérgéométriques et Hypérsphériques; Polynmes d'Hermite*, Gauthiers-Villars, Paris, 1926.

[4] Bailey, W. N. ; *Generalized Hypergeometric Series*, Cambridge Math. Tract No. 32, Cambridge University Press, Cambridge, 1935; Reprinted by Stechert-Hafner, New York, 1964.

[5] Brychkov, Yury A. ; *HAND BOOK OF Special Functions Derivatives, Integrals, Series and Other Formulas*, CRC press, Taylor & Francis Group, Boca Raton, London, New York, 2008.

[6] Choi, J., Harsh, H. and Rathie, A. K. ; Some Summation Formulas for the Appell's Function F_1 , *East Asian Math. J.* 17 (2001), 233–237.

- [7] Choi, J., Harsh, H. and Rathie, A. K. ; Further Summation Formulas for the Appell's Function F_1 , *J. Korea Soc. Math. Educ. Ser. B : Pure Appl. Math.* 12 (3) (2005), 223–228.
- [8] Choi, J., Rathie, A. K. and Malani, S. ; Kummer's Theorem and its Contiguous Identities, *Taiwanese Journal of Mathematics* 11 (5) (2007), 1521–1527.
- [9] Choi, J., Rathie, A. K. and Srivastava, H. M.; A Generalization of a Formula Due to Kummer, *Integral transforms and Special Functions*, 22 (11) (2011), 851–859.
- [10] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G. ; *Higher Transcendental Functions*, Vol. I, McGraw -Hill Book Company, New York, Toronto and London, 1953.
- [11] Gessel, I. and Stanton, D. ; Strange Evaluations of Hypergeometric Series, *SIAM. J. Math. Anal.*, 13(2), (1982), 295–308.
- [12] Heymann, W. ; Über Hypergeometrische Functionen, Deren Letztes ... Deren Letztis Element Speciell ist, *Z. Mathe. Phys.*, 44(1898), 3–44.
- [13] Heymann, W. ; Über Hypergeometrische Functionen Deren Letztes Element Speziell ist, *Z. Mathe. Phys.*, 44(1899), 280–288.
- [14] Karlsson, Per W. ; On two Hypergeometric Summation Formulas Conjectured by Gosper, *Simon Stevin*, 60(4), (1986), 329–337.
- [15] Kummer, E. E. ; Über die hypergeometrische Reihe

$$1 + \frac{\alpha, \beta}{1, \gamma} x + \frac{\alpha(\alpha + 1), \beta(\beta + 1)}{1, 2, \gamma(\gamma + 1)} x^2 + \frac{\alpha(\alpha + 1)(\alpha + 2), \beta(\beta + 1)(\beta + 2)}{1, 2, 3, \gamma(\gamma + 1)(\gamma + 2)} x^3 + \dots,$$
J. Reine Angew. Math. 15 (1836), 39–83 and 127–172; see also *Collected papers*, Vol. II: *Function Theory, Geometry and Miscellaneous* (Edited and with a Foreword by André Weil), Springer-Verlag, Berlin, Heidelberg and New York, 1975.
- [16] Lavoie, J. L. and Trottier, G. ; On the Sum of Certain Appell Series, *Ganita* 20 (1)(1969), 43–46.
- [17] Luke, Y. L. ; *Mathematical Functions and Their Approximations*, Academic Press, London, 1975.
- [18] Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O. I. ; *Integrals and Series, Volume III : More Special Functions*, Nauka, Moscow, 1986 (In Russian); (Translated from the Russian by G.G.Gould) Gordon and Breach Science Publishers, New York, 1990.
- [19] Qureshi, M. I. and Baboo, M. S. ; Some Unified and Generalized Kummer's First Summation Theorems with Applications in Laplace Transform Technique, *Asia Pac. J. Math.* 3 (1) (2016), 10–23.
- [20] Qureshi, M. I. and Baboo, M. S. ; Some Unified and Generalized Kummer's Second Summation Theorems with Applications in Laplace Transform Technique, *Int. J. Math. Appl.* 4(1-C) (2016) 45–52.
- [21] Qureshi, M. I. and Baboo, M. S. ; Some Unified and Generalized Kummer's Third Summation Theorems with Applications in Laplace Transform Technique, First International Conference cum Exhibition on Building Utility, Faculty of Engineering and Technology Jamia Millia Islamia, New Delhi, India, ICEBU, 1-3 December- 2016, pp.139–150. Enriched Publications Pvt.Ltd. New Delhi 110075 India.
- [22] Rainville, E. D. ; *Special Functions*, The Macmillan Company, New York, 1960 ; Reprinted by Chelsea Publ. Co., Bronx, New York, 1971.
- [23] Rakha, M. A. and Rathie, A. K. ; Generalizations of Classical Summation Theorems for the Series ${}_2F_1$ and ${}_3F_2$ with Applications, *Integr. Transforms Spec. Funct.* 22 (11) (2011), 823–840.
- [24] Shashikant, Sharma, S. and Rathie, A. K. ; Some Summation Formulas for the Appell's Function F_1 , *Proc. of 4th Int. Conf. SSFA.* 4 (2003), 81–84.
- [25] Slater, L. J. ; *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [26] Spiegel, M. R. ; Some Interesting Cases of the Hypergeometric Series, *Amer. Math. Monthly* 69 (1962), 894–896.
- [27] Srivastava, H. M. ; Hypergeometric Functions of Three Variables, *Ganita* 15 (2) (1964), 97–108.
- [28] Srivastava, H. M. and Manocha, H. L. ; *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1984.