

## LAPLACE TRANSFORMS OF MULTIPLE HYPERGEOMETRIC FUNCTIONS USING MELLIN-BARNES TYPE CONTOUR INTEGRATION

M.I. QURESHI<sup>1</sup>, DEEPAK KUMAR KABRA<sup>2,\*</sup> AND M. S. BABOO<sup>3</sup>

<sup>1</sup>Department of Applied Sciences and Humanities,  
Faculty of Engineering and Technology, Jamia Millia Islamia (A Central University), New Delhi - 110  
025, India

<sup>2</sup>Department of Mathematics, Govt. College Kota,  
University of Kota, Kota, Rajasthan- 324 001, India

<sup>3</sup>Department of Applied Sciences and Humanities,  
Knowledge park - III, Sharda University, Gr. Noida, U.P. - 301206, India

\*Corresponding author

**ABSTRACT.** In this paper, we find some results on Laplace transforms of Multiple hypergeometric functions like product theorems of Bailey, Cayley, Clausen, Orr, Preece, Watson for Gauss functions  ${}_2F_1$ , Appell's four functions, Ramanujan's product theorems, Henrici's triple product theorem, Whipple's quadratic transformation, Gauss, Goursat quadratic and cubic transformations, Product theorems of Kummer's functions  ${}_1F_1$ , Product theorems of Bessel's functions  ${}_0F_1$ , Bailey's cubic transformations et-cetera, in terms of Meijer's G-function of one variable and generalized hypergeometric function of one variable, using Mellin-Barnes type contour integral technique. The results presented here are presumably new.

2010 Mathematics Subject Classification. Primary:44A10; Secondary: 33C20, 33C60.

Key words and phrases. Multiple Hypergeometric functions, Laplace transforms, Meijer's G - function, Gauss's multiplication formula.

### 1. INTRODUCTION AND PRELIMINARIES

Throughout our present paper, we use the following standard notations:

$\mathbb{N} := \{1, 2, 3, \dots\}$ ,  $\mathbb{N}_0 := \{0, 1, 2, 3, \dots\} = \mathbb{N} \cup \{0\}$  and  $\mathbb{Z}^- := \{-1, -2, -3, \dots\} = \mathbb{Z}_0^- \setminus \{0\}$ .

Here, as usual,  $\mathbb{Z}$  denotes the set of integers,  $\mathbb{R}$  denotes the set of real numbers,  $\mathbb{R}^+$  denotes the set of positive real numbers and  $\mathbb{C}$  denotes the set of complex numbers.

The Pochhammer symbol (or the shifted factorial)  $(\lambda)_\nu$  ( $\lambda, \nu \in \mathbb{C}$ ) is defined, in terms of the familiar Gamma function, by

$$(1.1) \quad (\lambda)_\nu := \frac{\Gamma(\lambda + \nu)}{\Gamma(\lambda)} = \begin{cases} 1 & ; (\nu = 0; \lambda \in \mathbb{C} \setminus \{0\}) \\ \lambda(\lambda + 1) \dots (\lambda + n - 1) & ; (\nu = n \in \mathbb{N}; \lambda \in \mathbb{C}) \end{cases}$$

it is being understood *conventionally* that  $(0)_0 := 1$  and assumed tacitly that the Gamma quotient exists.

Of all the integrals which contain gamma functions in their integrands the most important ones are the so-called Mellin-Barnes integrals. Such integrals were first introduced by S. Pincherle in the year 1888; their theory has been developed by H. Mellin in the year 1910 and they were used for a complete integration of the hypergeometric differential equation by E.W. Barnes in the year 1908. The generalized Hypergeometric function [19, p. 100 Th. 35 and p. 102 Th. 36] is defined by means of Mellin-Barnes type contour integral in the following form, when  $p \leq q + 1$  then

$$(1.2) \quad \frac{\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)}{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)} {}_pF_q \left[ \alpha_1, \alpha_2, \dots, \alpha_p; \beta_1, \beta_2, \dots, \beta_q; z \right]$$

$$= \frac{1}{2\pi\omega} \int_{L_1} \frac{(-z)^\xi \Gamma(-\xi) \Gamma(\alpha_1 + \xi), \dots, \Gamma(\alpha_p + \xi)}{\Gamma(\beta_1 + \xi), \dots, \Gamma(\beta_q + \xi)} d\xi$$

where  $L_1$  is a suitable Mellin-Barnes path of integration [See 19, p. 95 figure (5), p. 98 figure(6)] and  $\alpha_i, \beta_j \in \mathbb{C} \setminus \mathbb{Z}_0^-, \{i = 1, 2, 3, \dots, p, j = 1, 2, 3, \dots, q\}$  and  $\omega = \sqrt{-1}, z \neq 0$ .

When  $p = q + 1$  then  $|\arg(-z)| < \pi$  and suppose that  $|z| < 1$ .

When  $p = q$  then  $|\arg(-z)| < \frac{\pi}{2}$  i.e.  $\Re(z) < 0$ .

When  $p < q$  then equation (1.2) is also valid.

Gauss's multiplication formula for the product of gamma functions [19, p.26 Th. 10; 23, p.23 (27)], is given by .

$$(1.3) \quad \Gamma(mz) = (2\pi)^{\frac{1-m}{2}} m^{(mz-\frac{1}{2})} \Gamma(z) \Gamma\left(z + \frac{1}{m}\right) \Gamma\left(z + \frac{2}{m}\right) \dots \Gamma\left(z + \frac{m-1}{m}\right)$$

where  $m$  is positive integer;  $mz \in \mathbb{C} \setminus \mathbb{Z}_0^-$ . For  $m = 2$  and  $m = 3$  we get Legendre's duplication formula and triplication formula respectively.

Using formula (1.3), we obtain

$$(1.4) \quad \Gamma(c + ks) = \Gamma\left\{k\left(s + \frac{c}{k}\right)\right\} = (2\pi)^{\frac{1-k}{2}} k^{(c-\frac{1}{2}+ks)} \prod_{q=1}^k \Gamma\left(s + \frac{c+q-1}{k}\right)$$

where  $k = 1, 2, 3, \dots$  and  $c + ks \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

In an attempt to give a meaning to the symbol  ${}_pF_q$ , when  $p > q + 1$ , Meijer introduced the G-function into Mathematical Analysis. Firstly the G-function was defined by Meijer [11] in the year 1936 by means of a finite series of generalized hypergeometric functions. Later on the Meijer's G-function of order  $(m, n, p, q)$  was defined by means of Mellin-Barnes type contour integral formula [5, p.207 (5.3.1); see also 9, p.143 (5.2.1);

10, p.2 (1.1.1, 1.1.3); 12, p.83; 13, p.1064 (21)], in the following form

$$(1.5) \quad G_{p,q}^{m,n} \left( z \left| \begin{array}{c} a_1, a_2, \dots, a_n; a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m; b_{m+1}, \dots, b_q \end{array} \right. \right) = \frac{1}{2\pi\omega} \int_{L_2} \frac{\prod_{j=1}^m \Gamma(b_j - s) \prod_{j=1}^n \Gamma(1 - a_j + s)}{\prod_{j=m+1}^q \Gamma(1 - b_j + s) \prod_{j=n+1}^p \Gamma(a_j - s)} z^s ds$$

( $a_k - b_j \neq 1, 2, 3, \dots$ ;  $k = 1, 2, 3, \dots, n$  and  $j = 1, 2, 3, \dots, m$ )

where  $0 \leq m \leq q, 0 \leq n \leq p$ ;  $z \neq 0$  and  $L_2$  is a suitable contour (See three cases of contour in the monographs [5, p.207 (2,3,4); 9, p.144 (2,3,4); 16, p.617 (1,2,3,4)]) and an empty product is interpreted as 1 and the parameters are such that no pole of  $\Gamma(b_j - s)$ ,  $j = 1, 2, 3, \dots, m$  coincides with any pole of  $\Gamma(1 - a_k + s)$ ,  $k = 1, 2, 3, \dots, n$ . Without any loss of generality, we are assuming that  $p \leq q$ . The MacRobert's E-function is the particular case of Meijer's G-function.

The G-function is symmetric with respect to order of the parameters in four groups  $a_1, a_2, \dots, a_n$ ;  $a_{n+1}, \dots, a_p$ ;  $b_1, b_2, \dots, b_m$ ;  $b_{m+1}, \dots, b_q$  individually.

If no pair among the parameters  $b_1, b_2, \dots, b_m$  may differ by an integer or zero (i.e. all poles are of the first order) then Meijer's function  $G_{p,q}^{m,n}(z)$  can be expressed as a sum of  $m$ -generalized hypergeometric functions  ${}_pF_{q-1}((-1)^{p-m-n}z)$  under the condition ( $p < q$  and for all finite values of  $z$ ) or ( $p = q$  and  $|z| < 1$ ).

If no pair among the parameters  $a_1, a_2, \dots, a_n$  may differ by an integer or zero (i.e. all poles are of the first order) then Meijer's function  $G_{p,q}^{m,n}(z)$  can be expressed as a sum of  $n$ -generalized hypergeometric functions  ${}_qF_{p-1}((-1)^{q-m-n}z^{-1})$  under the condition ( $p > q$  and for all finite values of  $z$ ) or ( $p = q$  and  $|z| > 1$ ).

If one (or more) pair among the parameters  $a_1, a_2, \dots, a_n$  or  $b_1, b_2, \dots, b_m$  may differ by an integer or zero then Logarithmic forms of the G-function occur due to the appearance of the poles of the higher order than unity, in the integrand of contour integral (1.5).

Property for cancellation of the numerator and denominator parameters

$$(1.6) \quad G_{p,q}^{m,n} \left( z \left| \begin{array}{c} a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m, b_{m+1}, \dots, b_{q-1}, a_1 \end{array} \right. \right) = G_{p-1,q-1}^{m,n-1} \left( z \left| \begin{array}{c} a_2, a_3, \dots, a_n, a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m, b_{m+1}, \dots, b_{q-1} \end{array} \right. \right)$$

where  $n, p, q \geq 1$ .

$$(1.7) \quad G_{p,q}^{m,n} \left( z \left| \begin{array}{c} a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_{p-1}, b_1 \\ b_1, b_2, \dots, b_m, b_{m+1}, \dots, b_q \end{array} \right. \right) = G_{p-1,q-1}^{m-1,n} \left( z \left| \begin{array}{c} a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_{p-1} \\ b_2, b_3, \dots, b_m, b_{m+1}, \dots, b_q \end{array} \right. \right)$$

where  $m, p, q \geq 1$ .

Translation property [13, p.1066 (24)]

$$\begin{aligned} & z^\sigma G_{p,q}^{m,n} \left( z \left| \begin{array}{c} a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m, b_{m+1}, \dots, b_q \end{array} \right. \right) \\ &= G_{p,q}^{m,n} \left( z \left| \begin{array}{c} a_1 + \sigma, a_2 + \sigma, \dots, a_n + \sigma, a_{n+1} + \sigma, \dots, a_p + \sigma \\ b_1 + \sigma, b_2 + \sigma, \dots, b_m + \sigma, b_{m+1} + \sigma, \dots, b_q + \sigma \end{array} \right. \right) \end{aligned}$$

Symmetric property (Transformation formula)

$$(1.8) \quad G_{p,q}^{m,n} \left( z \left| \begin{array}{c} a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m, b_{m+1}, \dots, b_q \end{array} \right. \right) \\ = G_{q,p}^{n,m} \left( \frac{1}{z} \left| \begin{array}{c} 1 - b_1, 1 - b_2, \dots, 1 - b_m, 1 - b_{m+1}, \dots, 1 - b_q \\ 1 - a_1, 1 - a_2, \dots, 1 - a_n, 1 - a_{n+1}, \dots, 1 - a_p \end{array} \right. \right)$$

Reduction formula between  ${}_pF_q$  and G-functions [5, p.215 (5.6.1); 23, p.47 (9)], is given by when  $p \leq q + 1$ , then

$$(1.9) \quad \frac{\Gamma(a_1)\Gamma(a_2)\dots\Gamma(a_p)}{\Gamma(b_1)\Gamma(b_2)\dots\Gamma(b_q)} {}_pF_q \left[ \begin{array}{c} a_1, a_2, \dots, a_p; \\ b_1, b_2, \dots, b_q; \end{array} z \right] = G_{p,q+1}^{1,p} \left( -z \left| \begin{array}{c} 1 - a_1, 1 - a_2, \dots, 1 - a_p \\ 0, 1 - b_1, 1 - b_2, \dots, 1 - b_q \end{array} \right. \right) \\ = G_{q+1,p}^{p,1} \left( -\frac{1}{z} \left| \begin{array}{c} 1, b_1, b_2, \dots, b_q \\ a_1, a_2, \dots, a_p \end{array} \right. \right) = (-z)^1 G_{p,q+1}^{1,p} \left( -z \left| \begin{array}{c} -a_1, -a_2, \dots, -a_p \\ -1, -b_1, -b_2, \dots, -b_q \end{array} \right. \right)$$

where ( $p \leq q$  and  $|z| < \infty$ ) or ( $p = q + 1$  and  $|z| < 1$ ).

For transformations and reduction formulas of multiple hypergeometric functions and product theorems of hypergeometric functions, we refer the monographs and research papers [1,2,5,7,8,9,16,19,21,23].

## 2. LAPLACE TRANSFORM

If there exists a number “ $M$ ” independent of  $t$  so that  $|\frac{f(t)}{g(t)}| \leq M$  as  $t \rightarrow t_0$  in the region  $R$  of the complex  $z$ -plane, where  $g(t) \neq 0$ , then we say that  $f(t) = O[g(t)]$  as  $t \rightarrow t_0$  in  $R$ .

If  $\lim_{t \rightarrow t_0} \frac{f(t)}{g(t)} = 0$ , then we say that  $f(t) = o[g(t)]$  as  $t \rightarrow t_0$  in  $R$ .

Let the function  $f(t)$  be piecewise continuous on the closed interval  $0 \leq t \leq T$  for every finite  $T > 0$ . Also let

$$(2.1) \quad f(t) = O[e^{\alpha t}], \quad t \rightarrow \infty$$

for some  $\alpha$ . Operational images (or operational representations) of many classes of special functions in the classical Laplace transform

$$(2.2) \quad \mathfrak{L}\{f(t) : p\} = \int_0^\infty e^{-pt} f(t) dt = F(p), \quad \Re(p) > \alpha,$$

can be obtained by appealing to Euler’s integral

$$(2.3) \quad \int_0^\infty e^{-pt} t^{\lambda-1} dt = \frac{\Gamma(\lambda)}{p^\lambda}$$

where  $\min\{\Re(\lambda), \Re(p)\} > 0$  or ( $\Re(p) = 0, 0 < \Re(\lambda) < 1$ ).

The integral (2.2) appeared for the first time in Euler’s investigation in the year 1737.

The regular use of the transformation of the form (2.2) began after the publication of P. S. Laplace's book in the year 1812. At the present time the Laplace transformation (2.2) is the most usable integral transformation. A complete account or elements of the theory of Laplace transformation can be found in numerous books on Laplace transformation, on operational calculus or on integral transformations. Among them we mention the monographs [3,4,6,14,15,17,18,20,22,24,25].

### 3. THEOREM ON LAPLACE TRANSFORMS

**Statement:**

When  $A \leq B+1$ ,  $\Re(p) > 0$ ,  $\Re(c) > 0$  and  $k$  is a positive integer. Suppose  $(a_A)$  abbreviates the array of  $A$  parameters given by  $a_1, a_2, \dots, a_A$  with similar interpretation for  $(b_B)$ , then

$$(3.1) \quad \mathfrak{L} \left\{ t^{c-1} {}_A F_B \left[ \begin{matrix} (a_A); \\ (b_B); \end{matrix} ; p \right] \right\} \\ = \frac{(2\pi)^{\frac{1-k}{2}} \prod_{j=1}^B \Gamma(b_j)}{p^c \prod_{i=1}^A \Gamma(a_i)} k^{(c-\frac{1}{2})} G_{1+B, A+k}^{A+k, 1} \left( \frac{-p^k}{yk^k} \left| \begin{matrix} 1, b_1, b_2, \dots, b_B \\ a_1, a_2, \dots, a_A, \frac{c}{k}, \frac{c+1}{k}, \dots, \frac{c+k-1}{k} \end{matrix} \right. \right)$$

$$(3.2) \quad \times G_{A+k, 1+B}^{1, A+k} \left( \frac{-yk^k}{p^k} \left| \begin{matrix} 1 - a_1, 1 - a_2, \dots, 1 - a_A, 1 + \left(\frac{-c}{k}\right), 1 + \left(\frac{-c-1}{k}\right), \dots, 1 + \left(\frac{-c-k+1}{k}\right) \\ 0, 1 - b_1, 1 - b_2, \dots, 1 - b_B \end{matrix} \right. \right)$$

provided that the right hand sides of (3.1) and (3.2) are convergent [See 5, p.207 (2,3,4); 9, p.144 (2,3,4); 16, p.617 (1,2,3,4)] and  $a_1, a_2, \dots, a_A, b_1, b_2, \dots, b_B \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Proof:** Consider the left hand side of equation (3.1):

$$\mathfrak{L} \left\{ t^{c-1} {}_A F_B \left[ \begin{matrix} (a_A); \\ (b_B); \end{matrix} ; p \right] \right\} = \int_0^\infty e^{-pt} t^{c-1} {}_A F_B \left[ \begin{matrix} (a_A); \\ (b_B); \end{matrix} ; p \right] dt$$

$$= \int_0^\infty e^{-pt} t^{c-1} \left( \frac{1}{2\pi\omega} \frac{\prod_{j=1}^B \Gamma(b_j)}{\prod_{i=1}^A \Gamma(a_i)} \int_L \frac{(-yt^k)^s \Gamma(-s) \prod_{i=1}^A \Gamma(a_i + s)}{\prod_{j=1}^B \Gamma(b_j + s)} ds \right) dt$$

where  $L$  is a suitable Mellin-Barnes type contour.

Now changing the order of integration, we get

$$\begin{aligned} \mathfrak{L} \left\{ t^{c-1} {}_A F_B \left[ \begin{matrix} (a_A); \\ (b_B); \end{matrix} yt^k : p \right] \right\} &= \frac{1}{2\pi\omega} \frac{\prod_{j=1}^B \Gamma(b_j)}{\prod_{i=1}^A \Gamma(a_i)} \int_L \left( \frac{(-y)^s \Gamma(-s) \prod_{i=1}^A \Gamma(a_i + s)}{\prod_{j=1}^B \Gamma(b_j + s)} \left( \int_0^\infty e^{-pt} t^{c+ks-1} dt \right) \right) ds \\ &= \frac{1}{p^c} \frac{1}{2\pi\omega} \frac{\prod_{j=1}^B \Gamma(b_j)}{\prod_{i=1}^A \Gamma(a_i)} \int_L \frac{\left( \frac{-y}{p^k} \right)^s \Gamma(-s) \Gamma(c + ks) \prod_{i=1}^A \Gamma(a_i + s)}{\prod_{j=1}^B \Gamma(b_j + s)} ds \end{aligned}$$

Now using Gauss's multiplication formula (1.4) for  $\Gamma(c + ks)$ , we get

$$\begin{aligned} \mathfrak{L} \left\{ t^{c-1} {}_A F_B \left[ \begin{matrix} (a_A); \\ (b_B); \end{matrix} yt^k : p \right] \right\} &= \frac{(2\pi)^{\frac{1-k}{2}} \prod_{j=1}^B \Gamma(b_j)}{p^c \prod_{i=1}^A \Gamma(a_i)} k^{(c-\frac{1}{2})} \times \\ &\times \frac{1}{2\pi\omega} \int_L \frac{\Gamma(-s) \prod_{i=1}^A \Gamma(a_i + s) \prod_{q=1}^k \Gamma\left(s + \frac{c+q-1}{k}\right)}{\prod_{j=1}^B \Gamma(b_j + s)} \left( \frac{-yk^k}{p^k} \right)^s ds \end{aligned}$$

Using definition (1.5) and transformation formula (1.8) of G-function, we get main results (3.2) and (3.1) respectively.

#### 4. LAPLACE TRANSFORMS OF MULTIPLE HYPERGEOMETRIC FUNCTIONS

For convenience, we shall use the notation  $\Delta(N; \lambda)$  for array of  $N$  parameters given by  $\frac{\lambda}{N}, \frac{\lambda+1}{N}, \frac{\lambda+2}{N}, \dots, \frac{\lambda+N-1}{N}$ . The following formulas hold for those suitable values of parameters for which gamma factors of the numerator and denominator are finite. The Laplace transforms of following Multiple hypergeometric functions are not found in the available literature on Laplace transforms [3,4,6,14,15,17,18,20,22,24,25]. Making

suitable adjustment of parameters and variables in equation (3.1), using the cancellation property (1.7), applying reduction formula (1.9), Legendre's duplication formula and triplication formula for the product of Gamma functions, after simplification we can find the following results, valid under the conditions associated with the result (3.1).

**Case(1):** Put  $c = 1, A = 3, B = 2, a_1 = a, a_2 = b, a_3 = c, b_1 = 1 + a - b, b_2 = 1 + a - c, y = 1, k = 1$ , in equation (3.1), we have

$$(4.1) \quad \mathfrak{L} \left\{ (1-t)^{-a} {}_3F_2 \left[ \begin{matrix} \frac{a}{2}, \frac{a+1}{2}, 1+a-b-c; \\ 1+a-b, 1+a-c; \end{matrix} -\frac{4t}{(1-t)^2} : p \right] \right\} \\ = \frac{\Gamma(1+a-b)\Gamma(1+a-c)}{p\Gamma(a)\Gamma(b)\Gamma(c)} G_{3,4}^{4,1} \left( -p \left| \begin{matrix} 1, 1+a-b, 1+a-c \\ a, b, c, 1 \end{matrix} \right. \right)$$

where  $1+a-b, 1+a-c, a, b, c \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  ${}_3F_2[.]$  is Clausenian hypergeometric functions.

**Case(2):** Put  $c = 1, A = 3, B = 2, a_1 = a, a_2 = 2b - a - 1, a_3 = a + 2 - 2b, b_1 = b, b_2 = a - b + \frac{3}{2}, y = \frac{1}{4}, k = 1$ , in equation (3.1), we have

$$(4.2) \quad \mathfrak{L} \left\{ (1-t)^{-a} {}_3F_2 \left[ \begin{matrix} \frac{a}{3}, \frac{a+1}{3}, \frac{a+2}{3}; \\ b, a-b+\frac{3}{2}; \end{matrix} -\frac{27t}{4(1-t)^3} : p \right] \right\} \\ = \frac{\Gamma(b)\Gamma(a-b+\frac{3}{2})}{p\Gamma(a)\Gamma(2b-a-1)\Gamma(a+2-2b)} G_{3,4}^{4,1} \left( -4p \left| \begin{matrix} 1, b, a-b+\frac{3}{2} \\ a, 2b-a-1, a+2-2b, 1 \end{matrix} \right. \right)$$

where  $a, b, a-b+\frac{3}{2}, 2b-a-1, a+2-2b \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(3):** Put  $c = 1, A = 3, B = 2, a_1 = a, a_2 = b - \frac{1}{2}, a_3 = 1 + a - b, b_1 = 2b, b_2 = 2 + 2a - 2b, y = 1, k = 1$ , in equation (3.1), we have

$$(4.3) \quad \mathfrak{L} \left\{ \left(1 - \frac{t}{4}\right)^{-a} {}_3F_2 \left[ \begin{matrix} \frac{a}{3}, \frac{a+1}{3}, \frac{a+2}{3}; \\ b, a-b+\frac{3}{2}; \end{matrix} \frac{27t^2}{(4-t)^3} : p \right] \right\} \\ = \frac{2^{(2a)} (b - \frac{1}{2}) \Gamma(b)\Gamma(a-b+\frac{3}{2})}{\pi p \Gamma(a)} G_{3,4}^{4,1} \left( -p \left| \begin{matrix} 1, 2b, 2+2a-2b \\ a, b-\frac{1}{2}, 1+a-b, 1 \end{matrix} \right. \right)$$

where  $a, b, a-b+\frac{3}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(4):** Put  $c = 1, A = 2, B = 1, a_1 = a, a_2 = b, b_1 = 2a, y = 1, k = 1$ , in equation (3.1), we have

$$(4.4) \quad \mathfrak{L} \left\{ \left(1 - \frac{t}{2}\right)^{-b} {}_2F_1 \left[ \begin{matrix} \frac{b}{2}, \frac{b+1}{2}; \\ a + \frac{1}{2}; \end{matrix} \left(\frac{t}{2-t}\right)^2 \right] : p \right\} = \frac{2^{(2a-1)} \Gamma(a + \frac{1}{2})}{\sqrt{\pi} p \Gamma(b)} G_{2,3}^{3,1} \left( -p \left| \begin{matrix} 1, 2a \\ a, b, 1 \end{matrix} \right. \right)$$

where  $a + \frac{1}{2}, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  ${}_2F_1[\cdot]$  is Gauss ordinary hypergeometric function.

**Case(5):** Put  $c = 1, A = 2, B = 1, a_1 = a, a_2 = a + \frac{1}{2} - b, b_1 = b + \frac{1}{2}, y = 1, k = 2$ , in result 1, we have

$$(4.5) \quad \mathfrak{L} \left\{ (1+t)^{-2a} {}_2F_1 \left[ \begin{matrix} a, b; \\ 2b; \end{matrix} \frac{4t}{(1+t)^2} \right] : p \right\} = \frac{\Gamma(b + \frac{1}{2})}{p \Gamma(a) \Gamma(a - b + \frac{1}{2}) \sqrt{\pi}} G_{2,4}^{4,1} \left( -\frac{p^2}{4} \left| \begin{matrix} 1, b + \frac{1}{2} \\ a, a - b + \frac{1}{2}, \frac{1}{2}, 1 \end{matrix} \right. \right)$$

where  $a - b + \frac{1}{2}, a, b + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(6):** Put  $c = 1, A = 2, B = 1, a_1 = 2a, a_2 = 2b, b_1 = a + b + \frac{1}{2}, y = 1, k = 1$ , in equation (3.1), we have

$$(4.6) \quad \mathfrak{L} \left\{ {}_2F_1 \left[ \begin{matrix} a, b; \\ a + b + \frac{1}{2}; \end{matrix} 4t(1-t) \right] : p \right\} = \frac{\Gamma(a + b + \frac{1}{2})}{\Gamma(2a)\Gamma(2b)p} G_{2,3}^{3,1} \left( -p \left| \begin{matrix} 1, a + b + \frac{1}{2} \\ 2a, 2b, 1 \end{matrix} \right. \right)$$

where  $a + b + \frac{1}{2}, 2a, 2b \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(7):** Put  $c = 1, A = 3, B = 2, a_1 = 2a, a_2 = 2b, a_3 = a + b, b_1 = 2a + 2b - 1, b_2 = a + b + \frac{1}{2}, y = 1, k = 1$ , in equation (3.1), we have

$$(4.7) \quad \mathfrak{L} \left\{ {}_2F_1 \left[ \begin{matrix} a, b; \\ a + b - \frac{1}{2}; \end{matrix} t \right] {}_2F_1 \left[ \begin{matrix} a, b; \\ a + b + \frac{1}{2}; \end{matrix} t \right] : p \right\} \\ = \frac{\sqrt{\pi} \Gamma(a + b - \frac{1}{2}) \Gamma(a + b + \frac{1}{2})}{p \Gamma(a) \Gamma(a + \frac{1}{2}) \Gamma(b) \Gamma(b + \frac{1}{2})} G_{3,4}^{4,1} \left( -p \left| \begin{matrix} 1, 2a + 2b - 1, a + b + \frac{1}{2} \\ 2a, 2b, a + b, 1 \end{matrix} \right. \right)$$

where  $a + b + \frac{1}{2}, a + b - \frac{1}{2}, a, b, a + \frac{1}{2}, b + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(8):** Put  $c = 1, A = 3, B = 2, a_1 = 2a, a_2 = 2b - 1, a_3 = a + b - 1, b_1 = 2a + 2b - 2, b_2 = a + b - \frac{1}{2}, y = 1, k = 1$ , in equation (3.1), we have

$$(4.8) \quad \mathfrak{L} \left\{ {}_2F_1 \left[ \begin{matrix} a, b; \\ a + b - \frac{1}{2}; \end{matrix} t \right] {}_2F_1 \left[ \begin{matrix} a, b - 1; \\ a + b - \frac{1}{2}; \end{matrix} t \right] : p \right\} \\ = \frac{\sqrt{\pi} \Gamma(a + b - \frac{1}{2}) \Gamma(a + b - \frac{1}{2})}{p \Gamma(a) \Gamma(a + \frac{1}{2}) \Gamma(b - \frac{1}{2}) \Gamma(b)} G_{3,4}^{4,1} \left( -p \left| \begin{matrix} 1, 2a + 2b - 2, a + b - \frac{1}{2} \\ 2a, 2b - 1, a + b - 1, 1 \end{matrix} \right. \right)$$

where  $a + b - \frac{1}{2}, a, b, a + \frac{1}{2}, b - \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(9):** Put  $c = 1, A = 3, B = 2, a_1 = a - b + \frac{1}{2}, a_2 = b - a + \frac{1}{2}, a_3 = \frac{1}{2}, b_1 = a + b + \frac{1}{2}, b_2 =$



$\frac{3}{2} - a - b, y = 1, k = 1$ , in equation (3.1), we have

$$(4.9) \quad \mathfrak{L} \left\{ {}_2F_1 \left[ \begin{matrix} a, & b; \\ a + b + \frac{1}{2}; \end{matrix} t \right] {}_2F_1 \left[ \begin{matrix} \frac{1}{2} - a, & \frac{1}{2} - b; \\ \frac{3}{2} - a - b & ; \end{matrix} t \right] : p \right\} \\ = \frac{\Gamma(\frac{3}{2} - a - b)\Gamma(a + b + \frac{1}{2})}{p\Gamma(b - a + \frac{1}{2})\Gamma(a - b + \frac{1}{2})\sqrt{\pi}} G_{3,4}^{4,1} \left( -p \left| \begin{matrix} 1, a + b + \frac{1}{2}, \frac{3}{2} - a - b \\ a - b + \frac{1}{2}, b - a + \frac{1}{2}, \frac{1}{2}, 1 \end{matrix} \right. \right)$$

where  $a + b + \frac{1}{2}, \frac{3}{2} - a - b, b - a + \frac{1}{2}, a - b + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(10):** Put  $c = 1, A = 3, B = 2, a_1 = 2a, a_2 = 2b, a_3 = a + b, b_1 = 2a + 2b, b_2 = a + b + \frac{1}{2}, y = 1, k = 1$ , in equation (3.1), we have

$$(4.10) \quad \mathfrak{L} \left\{ \left( {}_2F_1 \left[ \begin{matrix} a, & b; \\ a + b + \frac{1}{2}; \end{matrix} t \right] \right)^2 : p \right\} = \frac{2\sqrt{\pi}\Gamma(a + b + \frac{1}{2})\Gamma(a + b + \frac{1}{2})}{p\Gamma(a)\Gamma(a + \frac{1}{2})\Gamma(b)\Gamma(b + \frac{1}{2})} G_{3,4}^{4,1} \left( -p \left| \begin{matrix} 1, 2a + 2b, a + b + \frac{1}{2} \\ 2a, 2b, a + b, 1 \end{matrix} \right. \right)$$

where  $a + b + \frac{1}{2}, a, b, a + \frac{1}{2}, b + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(11):** Put  $c = 1, A = 2, B = 1, a_1 = a, a_2 = b + c, b_1 = d, y = 1, k = 1$ , in equation (3.1), we have

$$(4.11) \quad \mathfrak{L} \{ \mathbf{F}_1[a; b, c; d; t, t] : p \} = \frac{\Gamma(d)}{p\Gamma(a)\Gamma(b + c)} G_{2,3}^{3,1} \left( -p \left| \begin{matrix} 1, d \\ a, b + c, 1 \end{matrix} \right. \right)$$

where  $d, a, b + c \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  $\mathbf{F}_1[\cdot]$  is Appell's function of first kind [23, p.53 (1.6.4)].

**Case(12):** Put  $c = 1, A = 3, B = 2, a_1 = \frac{a}{2}, a_2 = \frac{a + 1}{2}, a_3 = b, b_1 = \frac{c}{2}, b_2 = \frac{c + 1}{2}, y = 1, k = 2$ , in equation (3.1), we have

$$(4.12) \quad \mathfrak{L} \{ \mathbf{F}_1[a; b, b; c; t, -t] : p \} = \frac{2^{(a-c)}\Gamma(c)}{\sqrt{\pi}p\Gamma(a)\Gamma(b)} G_{3,5}^{5,1} \left( -\frac{p^2}{4} \left| \begin{matrix} 1, \Delta(2; c) \\ \Delta(2; a), b, \frac{1}{2}, 1 \end{matrix} \right. \right)$$

where  $c, a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(13):** Put  $c = 1, A = 4, B = 3, a_1 = \frac{a}{2}, a_2 = \frac{a + 1}{2}, a_3 = b, a_4 = c - b, b_1 = c, b_2 = \frac{c}{2}, b_3 = \frac{c + 1}{2}, y = 1, k = 2$ , in equation (3.1), we have

$$(4.13) \quad \mathfrak{L} \{ \mathbf{F}_2[a; b, b; c, c; t, -t] : p \} = \frac{2^{(a-c)}[\Gamma(c)]^2}{\sqrt{\pi}p\Gamma(a)\Gamma(b)\Gamma(c - b)} G_{4,6}^{6,1} \left( -\frac{p^2}{4} \left| \begin{matrix} 1, c, \Delta(2; c) \\ \Delta(2; a), b, c - b, \frac{1}{2}, 1 \end{matrix} \right. \right)$$

where  $c, a, b, c - b \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  $\mathbf{F}_2[\cdot]$  is Appell's function of second kind [23, p.53 (1.6.5)].

**Case(14):** Put  $c = 1, A = 4, B = 3, a_1 = \frac{a}{2}, a_2 = \frac{a + 1}{2}, a_3 = \frac{\lambda + \mu}{2}, a_4 = \frac{\lambda + \mu + 1}{2}, b_1 = \lambda + \mu, b_2 = \lambda + \frac{1}{2}, b_3 = \mu + \frac{1}{2}, y = 1, k = 2$ , in equation (3.1), we have

$$\mathfrak{L} \{ \mathbf{F}_2[a; \lambda, \mu; 2\lambda, 2\mu; t, -t] : p \} = \frac{2^{(\lambda + \mu + a - 2)}\Gamma(\lambda + \frac{1}{2})\Gamma(\mu + \frac{1}{2})}{\pi^{\frac{3}{2}}p\Gamma(a)} \times$$

$$(4.14) \quad \times G_{4,6}^{6,1} \left( -\frac{p^2}{4} \left| \begin{array}{c} 1, \lambda + \mu, \lambda + \frac{1}{2}, \mu + \frac{1}{2} \\ \Delta(2; a), \Delta(2; \lambda + \mu), \frac{1}{2}, 1 \end{array} \right. \right)$$

where  $a, \lambda + \frac{1}{2}, \mu + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(15):** Put  $c = 1, A = 4, B = 3, a_1 = a, a_2 = b, a_3 = \frac{a+b}{2}, a_4 = \frac{a+b+1}{2}, b_1 = a+b, b_2 = \frac{c}{2}, b_3 = \frac{c+1}{2}, y = 1, k = 2$ , in equation (3.1), we have

$$(4.15) \quad \mathfrak{L} \{ \mathbf{F}_3[a, a; b, b; c; t, -t] : p \} = \frac{2^{(a+b-c)} \Gamma(c)}{\sqrt{\pi} p \Gamma(a) \Gamma(b)} G_{4,6}^{6,1} \left( -\frac{p^2}{4} \left| \begin{array}{c} 1, a+b, \Delta(2; c) \\ a, b, \Delta(2; a+b), \frac{1}{2}, 1 \end{array} \right. \right)$$

where  $c, a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  $\mathbf{F}_3[.]$  is Appell's function of third kind [23, p.53 (1.6.6)].

**Case(16):** Put  $c = 1, A = 4, B = 3, a_1 = a, a_2 = b, a_3 = \frac{c+d}{2}, a_4 = \frac{c+d-1}{2}, b_1 = c, b_2 = d, b_3 = c+d-1, y = 4, k = 1$ , in equation (3.1), we have

$$(4.16) \quad \mathfrak{L} \{ \mathbf{F}_4[a, b; c, d; t, t] : p \} = \frac{2^{(c+d-2)} \Gamma(c) \Gamma(d)}{\sqrt{\pi} p \Gamma(a) \Gamma(b)} G_{4,5}^{5,1} \left( -\frac{p}{4} \left| \begin{array}{c} 1, c, d, c+d-1 \\ a, b, \Delta(2; c+d-1), 1 \end{array} \right. \right)$$

where  $c, d, a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  $\mathbf{F}_4[.]$  is Appell's function of fourth kind [23, p.53 (1.6.7)].

**Case(17):** Put  $c = 1, A = 4, B = 3, a_1 = \frac{a}{2}, a_2 = \frac{a+1}{2}, a_3 = \frac{b}{2}, a_4 = \frac{b+1}{2}, b_1 = c, b_2 = \frac{c}{2}, b_3 = \frac{c+1}{2}, y = -4, k = 2$ , in equation (3.1), we have

$$(4.17) \quad \mathfrak{L} \{ \mathbf{F}_4[a, b; c, c; t, -t] : p \} = \frac{2^{(a+b-c-1)} [\Gamma(c)]^2}{\pi p \Gamma(a) \Gamma(b)} G_{4,6}^{6,1} \left( \frac{p^2}{16} \left| \begin{array}{c} 1, c, \Delta(2; c) \\ \Delta(2; a), \Delta(2; b), \frac{1}{2}, 1 \end{array} \right. \right)$$

where  $c, a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(18):** Put  $c = 1, A = 1, B = 1, a_1 = a, b_1 = b, y = 1, k = 1$ , in equation (3.1), we have

$$(4.18) \quad \mathfrak{L} \left\{ e^t {}_1F_1 \left[ \begin{array}{c} b-a; \\ b \end{array} ; -t \right] : p \right\} = \frac{\Gamma(b)}{p \Gamma(a)} G_{2,2}^{2,1} \left( -p \left| \begin{array}{c} 1, b \\ a, 1 \end{array} \right. \right) \\ = \frac{1}{p} {}_2F_1 \left[ \begin{array}{c} a, 1; \\ b \end{array} ; \frac{1}{p} \right]$$

where  $a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  ${}_1F_1[.]$  is Kummer's confluent hypergeometric function.

**Case(19):** Put  $c = 1, A = 1, B = 1, a_1 = a, b_1 = 2a, y = 2, k = 1$ , in equation (3.1), we have

$$(4.19) \quad \mathfrak{L} \left\{ e^t {}_0F_1 \left[ \begin{array}{c} - \\ a + \frac{1}{2}; \end{array} ; \frac{t^2}{4} \right] : p \right\} = \frac{2^{(2a-1)} \Gamma(a + \frac{1}{2})}{p \sqrt{\pi}} G_{2,2}^{2,1} \left( -\frac{p}{2} \left| \begin{array}{c} 1, 2a \\ a, 1 \end{array} \right. \right) \\ = \frac{1}{p} {}_2F_1 \left[ \begin{array}{c} a, 1; \\ 2a \end{array} ; \frac{2}{p} \right]$$

where  $a + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(20):** Put  $c = 1, A = 2, B = 3, a_1 = a, a_2 = b - a, b_1 = b, b_2 = \frac{b}{2}, b_3 = \frac{b+1}{2}, y = \frac{1}{4}, k = 2$ , in equation (3.1), we have

$$(4.20) \quad \mathfrak{L} \left\{ {}_1F_1 \left[ \begin{matrix} a; \\ b; \end{matrix} t \right] {}_1F_1 \left[ \begin{matrix} a; \\ b; \end{matrix} -t \right] : p \right\} = \frac{[\Gamma(b)]^2}{2^{(b-1)} p \Gamma(a) \Gamma(b-a)} G_{4,4}^{4,1} \left( -p^2 \left| \begin{matrix} 1, b, \Delta(2; b) \\ a, b-a, \frac{1}{2}, 1 \end{matrix} \right. \right)$$

$$= \frac{1}{p} {}_4F_3 \left[ \begin{matrix} a, b-a, \frac{1}{2}, 1; \\ b, \Delta(2; b); \end{matrix} \frac{1}{p^2} \right]$$

where  $a, b, b-a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(21):** Put  $c = 1, A = 3, B = 8, a_1 = \frac{a+b-1}{2}, a_2 = \frac{a+b}{3}, a_3 = \frac{a+b+1}{3}, b_1 = a, b_2 = b, b_3 = \frac{a}{2}, b_4 = \frac{b}{2}, b_5 = \frac{a+1}{2}, b_6 = \frac{b+1}{2}, b_7 = \frac{a+b-1}{2}, b_8 = \frac{a+b}{2}, y = -\frac{27}{64}, k = 2$ , in equation (3.1), we have

$$(4.21) \quad \mathfrak{L} \left\{ {}_0F_2 \left[ \begin{matrix} -; \\ a, b; \end{matrix} t \right] {}_0F_2 \left[ \begin{matrix} -; \\ a, b; \end{matrix} -t \right] : p \right\} = \frac{3^{(a+b-\frac{3}{2})} [\Gamma(a)]^2 [\Gamma(b)]^2}{p 2^{(2a+2b-3)}} \times$$

$$\times G_{9,5}^{5,1} \left( \frac{16p^2}{27} \left| \begin{matrix} 1, a, b, \Delta(2; a), \Delta(2; b), \Delta(2; a+b-1) \\ \Delta(3; a+b-1), \frac{1}{2}, 1 \end{matrix} \right. \right)$$

$$= \frac{1}{p} {}_5F_8 \left[ \begin{matrix} \Delta(3; a+b-1), \frac{1}{2}, 1; \\ a, b, \Delta(2; a), \Delta(2; b), \Delta(2; a+b-1); \end{matrix} -\frac{27}{16p^2} \right]$$

where  $a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(22):** Put  $c = 1, A = 2, B = 3, a_1 = \frac{a+b}{2}, a_2 = \frac{a+b+1}{2}, b_1 = a + \frac{1}{2}, b_2 = b + \frac{1}{2}, b_3 = a+b, y = \frac{1}{4}, k = 2$ , in equation (3.1), we have

$$(4.22) \quad \mathfrak{L} \left\{ {}_1F_1 \left[ \begin{matrix} a; \\ 2a; \end{matrix} t \right] {}_1F_1 \left[ \begin{matrix} b; \\ 2b; \end{matrix} -t \right] : p \right\} = \frac{2^{(a+b-1)} \Gamma(a + \frac{1}{2}) \Gamma(b + \frac{1}{2})}{\pi p} \times$$

$$\times G_{4,4}^{4,1} \left( -p^2 \left| \begin{matrix} 1, a + \frac{1}{2}, b + \frac{1}{2}, a+b \\ \Delta(2; a+b), \frac{1}{2}, 1 \end{matrix} \right. \right)$$

$$= \frac{1}{p} {}_4F_3 \left[ \begin{matrix} \Delta(2; a+b), \frac{1}{2}, 1; \\ a + \frac{1}{2}, b + \frac{1}{2}, a+b; \end{matrix} \frac{1}{p^2} \right]$$

where  $a + \frac{1}{2}, b + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

**Case(23):** Put  $c = 1, A = 2, B = 7, a_1 = 3b - \frac{1}{4}, a_2 = 3b + \frac{1}{4}, b_1 = 6b, b_2 = 2b, b_3 =$

$2b + \frac{1}{3}, b_4 = 2b + \frac{2}{3}, b_5 = 4b - \frac{1}{3}, b_6 = 4b, b_7 = 4b + \frac{1}{3}, y = \frac{64}{729}, k = 3$ , in equation (3.1), we have

$$\begin{aligned}
(4.23) \quad \mathfrak{L} \left\{ {}_0F_1 \left[ \begin{matrix} -; \\ 6b; \end{matrix} t \right] {}_0F_1 \left[ \begin{matrix} -; \\ 6b; \end{matrix} \omega t \right] {}_0F_1 \left[ \begin{matrix} -; \\ 6b; \end{matrix} \omega^2 t \right] : p \right\} &= \frac{2^{(18b - \frac{5}{2})} [\Gamma(6b)]^3}{3^{(18b - \frac{5}{2})} p} \times \\
&\times G_{8,5}^{5,1} \left( -\frac{27p^3}{64} \left| \begin{matrix} 1, 6b, \Delta(3; 6b), \Delta(3; 12b - 1) \\ \Delta(2; 6b - \frac{1}{2}), \Delta(3; 1) \end{matrix} \right. \right) \\
&= \frac{1}{p} {}_5F_7 \left[ \begin{matrix} \Delta(2; 6b - \frac{1}{2}), \Delta(3; 1); \frac{64}{27p^3} \\ 6b, \Delta(3; 6b), \Delta(3; 12b - 1); \end{matrix} \right]
\end{aligned}$$

where  $\omega$  is the cube root of unity ( $\omega = e^{\frac{2\pi i}{3}}$ ) and  $6b \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

Case(24): Put  $c = 1, A = 0, B = 3, b_1 = a, b_2 = \frac{a}{2}, b_3 = \frac{a+1}{2}, y = -\frac{1}{4}, k = 2$ , in equation (3.1), we have

$$\begin{aligned}
(4.24) \quad \mathfrak{L} \left\{ {}_0F_1 \left[ \begin{matrix} -; \\ a; \end{matrix} t \right] {}_0F_1 \left[ \begin{matrix} -; \\ a; \end{matrix} -t \right] : p \right\} &= \frac{[\Gamma(a)]^2}{p 2^{(a-1)}} G_{4,2}^{2,1} \left( p^2 \left| \begin{matrix} 1, a, \Delta(2; a) \\ \frac{1}{2}, 1 \end{matrix} \right. \right) \\
&= \frac{1}{p} {}_2F_3 \left[ \begin{matrix} \frac{1}{2}, 1; -\frac{1}{p^2} \\ a, \Delta(2; a); \end{matrix} \right]
\end{aligned}$$

where  $a \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

Case(25): Put  $c = 1, A = 2, B = 3, a_1 = \frac{a+b}{2}, a_2 = \frac{a+b-1}{2}, b_1 = a, b_2 = b, b_3 = a+b-1, y = 4, k = 1$ , in equation (3.1), we have

$$\begin{aligned}
(4.25) \quad \mathfrak{L} \left\{ {}_0F_1 \left[ \begin{matrix} -; \\ a; \end{matrix} t \right] {}_0F_1 \left[ \begin{matrix} -; \\ b; \end{matrix} t \right] : p \right\} &= \frac{2^{(a+b-2)} \Gamma(a)\Gamma(b)}{\sqrt{\pi} p} G_{4,3}^{3,1} \left( -\frac{p}{4} \left| \begin{matrix} 1, a, b, a+b-1 \\ \Delta(2; a+b-1), 1 \end{matrix} \right. \right) \\
&= \frac{1}{p} {}_3F_3 \left[ \begin{matrix} \Delta(2; a+b-1), 1; \frac{4}{p} \\ a, b, a+b-1; \end{matrix} \right]
\end{aligned}$$

where  $a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

Case(26): Put  $c = 1, A = 1, B = 2, a_1 = a, b_1 = a + \frac{1}{2}, b_2 = 2a, y = \frac{1}{4}, k = 2$ , in equation (3.1), we have

$$\begin{aligned}
(4.26) \quad \mathfrak{L} \left\{ \left( {}_1F_1 \left[ \begin{matrix} a; \\ 2a; \end{matrix} t \right] \right)^2 : p \right\} &= \frac{[\Gamma(a + \frac{1}{2})]^2 2^{(2a-1)}}{\pi (p-1)} G_{3,3}^{3,1} \left( -(p-1)^2 \left| \begin{matrix} 1, a + \frac{1}{2}, 2a \\ a, \frac{1}{2}, 1 \end{matrix} \right. \right) \\
&= \frac{1}{(p-1)} {}_3F_2 \left[ \begin{matrix} a, \frac{1}{2}, 1; \frac{1}{(p-1)^2} \\ 2a, a + \frac{1}{2}; \end{matrix} \right]
\end{aligned}$$

where  $a + \frac{1}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-$  and  $\Re(p) > 1$ .

From case 1 to case 25 the condition of validity is  $\Re(p) > 0$ .

We conclude our present investigation by observing that several other consequences of Laplace transforms can also be deduced in an analogous manner.

#### REFERENCES

- [1] Andrews, G. E., Askey, R. and Roy, R.; *Special Function, Encyclopedia of Mathematics and its Applications*, Vol. 71, Cambridge University Press, Cambridge, 1999.
- [2] Bailey, W. N. ; *Generalized Hypergeometric Series*, Cambridge Math. Tract No. 32, Cambridge University Press, Cambridge, 1935; Reprinted by Stechert-Hafner, New York, 1964.
- [3] Cossar, J. and Erdélyi, A.; *Dictionary of Laplace Transforms*, Admiralty Computing Service, London, 1944 – 46.
- [4] Doetsch, G.; *Introduction to the Theory and Application of the Laplace Transformation*, Springer-Verlag, Berlin, 1974.
- [5] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F.G.; *Higher Transcendental Functions*, Vol. I, (Bateman Manuscript Project), McGraw-Hill Book Co. Inc., New York, Toronto and London, 1953.
- [6] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G.; *Tables of Integral Transforms*, Vol.I (Bateman Manuscript Project), McGraw-Hill Book Co. Inc., New York, Toronto and London, 1954.
- [7] Henrici, P.; A triple product theorem for hypergeometric series, *SIAM J. Math. Anal.*, 18(1987), 1513-1518.
- [8] Karlsson, Per W. and Srivastava, H. M.; A note on Henrici's triple product theorem, *Proc. Amer. Math. Soc.*, 110(1)(1990), 85-88.
- [9] Luke, Y. L.; *The Special Functions and Their Approximations*, Vol. 1, Academic Press, New York, 1969.
- [10] Mathai, A. M. and Saxena, R. K.; *Generalized Hypergeometric Functions with Applications in Statistics and Physical Sciences, Lectures Notes in Mathematics No. 348*, Springer-Verlag, Berlin, Heidelberg, New York, 1973.
- [11] Meijer, C. S.; Über Whittakersche Bezv. Besselsche Funktionen und deren Produkte, *Nieuw. Arch. Wiskunde*, 18(2)(1936), 10-39.
- [12] Meijer, C. S.; Neue Integralstellungen für Whittakersche Funktionen I, *Nederl. Akad. Wetensch. Proc. Ser. A*, 44(1941), 81-92.
- [13] Meijer, C. S.; Multiplikationstheoreme für die Funktion  $G_{p,q}^{m,n}(z)$ , *Nederl. Akad. Wetensch. Proc. Ser. A*, 44(1941), 1062-1070.
- [14] Nixon, F. E.; *Handbook of Laplace Transformation*, Prentice Hall, 1960.
- [15] Oberhettinger, F. and Badii, L.; *Tables of Laplace Transforms*, Springer-Verlag, New York, Heidelberg, Berlin, 1973.
- [16] Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O. I.; *Integrals and Series, Vol.3: More Special Functions*, Gordon and Breach Science Publishers, New York, 1990.
- [17] Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O. I.; *Integrals and Series, Vol.4: Direct Laplace Transform*, Gordon and Breach Science Publishers, New York, 1992.

- [18] Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O. I.; *Integrals and Series, Vol.5: Inverse Laplace Transform*, Gordon and Breach Science Publishers, New York, 1992.
- [19] Rainville, E. D.; *Special Functions*, The Macmillan Co. Inc., New York 1960; Reprinted by Chelsea Publ. Co. Bronx, New York, 1971.
- [20] Roberts, G. E. and Kaufman, H.; *Tables of Laplace Transforms*, W. B. Saunders Co., McAinsh Toronto, 1966.
- [21] Slater, L. J.; *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [22] Spiegel, M. R.; *Theory and Problem of Laplace Transforms*, Schaum's Outline Series, 1986.
- [23] Srivastava, H. M. and Manocha, H. L.; *A Treatise on Generating functions*, Halsted Press (Ellis Horwood Limited, Chichester, U. K.) John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1984.
- [24] Thompson, W. T.; *Laplace Transformation*, Prentice Hall, New York, 1950.
- [25] Widder, D. V.; *The Laplace Transform*, Princeton University Press, Princeton, New Jersey, 1946.