

**SUMMATION THEOREMS FOR ${}_p F_q[A \pm M, (\alpha_p); (\beta_q); Z]$ VIA
MELLIN-BARNES TYPE CONTOUR INTEGRAL AND ITS
APPLICATIONS**

M. I. QURESHI¹ AND M. S. BABOO^{2,*}

¹Department of Applied Sciences and Humanities, Faculty of Engineering and Technology,
Jamia Millia Islamia (A Central University), New Delhi -110025, India

²School of Basic Sciences and Research, Sharda University,
Greater Noida, Uttar Pradesh, 201306, India

*Corresponding author: mesub007@gmail.com

Received March 7, 2017

ABSTRACT. In this paper, we find summation theorems for generalized hypergeometric functions ${}_p F_q[a \pm m, (\alpha_p); (\beta_q); z]$ by means of Mellin-Barnes type contour integral representation. Some closed forms (believed to be new) of summation theorems for Clausen hypergeometric functions ${}_3F_2[a + m, b, c; a, 1 + b - c \pm p; -1]$, ${}_3F_2[a + m, b, c; a, b - c \pm p; -1]$, ${}_3F_2[a + m, b, c; a, (1 + b + c \pm p)/2; 1/2]$, ${}_3F_2[a + m, b, c; a, (b + c \pm p)/2; 1/2]$, ${}_3F_2[a + m, b, 1 - b \pm p; a, h; 1/2]$, ${}_3F_2[a + m, b, -b \pm p; a, h; 1/2]$, ${}_3F_2[a + m, b, c; a, c + 1; 1]$ and Gauss hypergeometric functions ${}_2F_1[a + m, b; 1 + a - b \pm p; -1]$, ${}_2F_1[a + m, b; a - b \pm p; -1]$, ${}_2F_1[a + m, b; (1 + a + b \pm p)/2; 1/2]$, ${}_2F_1[a + m, b; (a + b \pm p)/2; 1/2]$, ${}_2F_1[a + m, 1 - a \pm p; g; 1/2]$, ${}_2F_1[a + m, -a \pm p; g; 1/2]$ with suitable convergence conditions, are also obtained.

2010 Mathematics Subject Classification. Primary 33B15, 33C20; Secondary 33C05, 33C60.

Key words and phrases. generalized hypergeometric function; Legendre duplication formula; Mellin-Barnes type contour integral; Meijer's G-function; generalized Kummer's first, second and third summation theorems.

1. INTRODUCTION, DEFINITIONS AND PRELIMINARIES

In the usual notation, let \mathbb{R} and \mathbb{C} denote the sets of real and complex numbers, respectively. Also let

$$\mathbb{N}_0 = \mathbb{N} \cup \{0\}, \quad \mathbb{N} = \{1, 2, 3, \dots\} = \mathbb{N}_0 \setminus \{0\},$$

$$\mathbb{Z}_0^- = \{0, -1, -2, \dots\} = \mathbb{Z}^- \cup \{0\}, \quad \mathbb{Z}^- = \{-1, -2, -3, \dots\}$$

and $\mathbb{Z} = \mathbb{Z}_0^- \cup \mathbb{N}$ being the sets of integers.

In terms of Gamma function $\Gamma(z)$, the widely-used Pochhammer symbol $(\lambda)_\nu$ ($\lambda, \nu \in \mathbb{C}$) is defined, in general, by

$$(1.1) \quad (\lambda)_\nu := \frac{\Gamma(\lambda + \nu)}{\Gamma(\lambda)} = \begin{cases} 1, & (\nu = 0; \lambda \in \mathbb{C} \setminus \{0\}) \\ \lambda(\lambda + 1) \dots (\lambda + n - 1), & (\nu = n \in \mathbb{N}; \lambda \in \mathbb{C}), \end{cases}$$

it is being understood *conventionally* that $(0)_0 := 1$ and assumed *tacitly* that the Γ quotient exists (see, for details, [15])

$$(1.2) \quad \int_0^\infty e^{-st} t^{\alpha-1} dt = \frac{\Gamma(\alpha)}{s^\alpha},$$

$$\left(\Re(s) > 0, 0 < \Re(\alpha) < \infty, \text{ or } \Re(s) = 0, 0 < \Re(\alpha) < 1 \right).$$

Here *generalized hypergeometric function* ${}_pF_q$ with p numerator parameters $\alpha_1, \alpha_2, \dots, \alpha_p$ and q denominator parameters $\beta_1, \beta_2, \dots, \beta_q$ is defined by

$$(1.3) \quad {}_pF_q \left[\begin{matrix} \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{matrix} z \right] = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^p (\alpha_j)_n}{\prod_{j=1}^q (\beta_j)_n} \frac{z^n}{n!},$$

$$\left(p, q \in \mathbb{N}_0; p \leq q + 1; p \leq q \text{ and } |z| < \infty, \text{ or } p = q + 1 \text{ and } |z| < 1, \text{ or } p = q + 1, |z| = 1 \text{ and } \Re(\omega) > 0, \text{ or } p = q + 1, |z| = 1, z \neq 1 \text{ and } 0 \geq \Re(\omega) > -1 \right),$$

where

$$\omega := \sum_{j=1}^q \beta_j - \sum_{j=1}^p \alpha_j,$$

$$\left(\alpha_j \in \mathbb{C} (j = 1, 2, \dots, p); \beta_j \in \mathbb{C} \setminus \mathbb{Z}_0^- (j = 1, 2, \dots, q) \right).$$

Of all the integrals which contain gamma functions in their integrands the most important ones are the so-called Mellin-Barnes integrals. Such integrals were first introduced by S. Pincherle in the year 1888; their theory has been developed by H. Mellin in the year 1910 and they were used for a complete integration of the hypergeometric differential equation by E.W. Barnes in the year 1908. The generalized Hypergeometric function [13, p.100, Theorem

35 and p.102, Theorem 36] is defined by means of Mellin-Barnes type contour integral in the following form; when $p \leq q + 1$, then

$$(1.4) \quad \frac{\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)}{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)} {}_pF_q \left[\begin{matrix} \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{matrix} z \right] = \\ = \frac{1}{2\pi\omega} \int_{L_1} \frac{(-z)^\xi \Gamma(-\xi) \Gamma(\alpha_1 + \xi), \dots, \Gamma(\alpha_p + \xi)}{\Gamma(\beta_1 + \xi), \dots, \Gamma(\beta_q + \xi)} d\xi,$$

where L_1 is a suitable Mellin-Barnes path of integration see [13, p.95, figure (5), p.98, figure(6)] and $\alpha_i, \beta_j \in \mathbb{C} \setminus \mathbb{Z}_0^-, \{i = 1, 2, 3, \dots, p; j = 1, 2, 3, \dots, q\}$ and $\omega = \sqrt{-1}, z \neq 0$.

When $p = q + 1$ then $|\arg(-z)| < \pi$ and suppose that $|z| < 1$.

When $p = q$ then $|\arg(-z)| < \frac{\pi}{2}$ i.e. $\Re(z) < 0$.

When $p < q$ then equation (1.4) is also valid.

In an attempt to give a meaning to the symbol ${}_pF_q$, when $p > q + 1$. Firstly the G-function was defined by Meijer [6] in the year 1936 by means of a finite series of generalized hypergeometric functions. Later on the Meijer's G-function of order (m, n, p, q) was defined by means of Mellin-Barnes type contour integral formula [3, p.207, Entry (5.3.1)]; see also [4, p.143, Entry (5.2.1)]; [5, p.2, Entries (1.1.1, 1.1.3)] and [7, p.1064, Entry (21)], in the following form

$$(1.5) \quad G_{p,q}^{m,n} \left(z \left| \begin{matrix} a_1, a_2, \dots, a_n; a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m; b_{m+1}, \dots, b_q \end{matrix} \right. \right) = \\ = \frac{1}{2\pi\omega} \int_{L_2} \frac{\prod_{j=1}^m \Gamma(b_j - s) \prod_{j=1}^n \Gamma(1 - a_j + s)}{\prod_{j=m+1}^q \Gamma(1 - b_j + s) \prod_{j=n+1}^p \Gamma(a_j - s)} z^s ds,$$

$$(a_k - b_j \neq 1, 2, 3, \dots; k = 1, 2, 3, \dots, n \text{ and } j = 1, 2, 3, \dots, m),$$

where $0 \leq m \leq q, 0 \leq n \leq p; z \neq 0$ and L_2 is a suitable contour (See three cases of contour in the monographs [3, p.207, Equations (2,3,4)]; [4, p.144, Equations (2,3,4)]; [9, p.617, Equations (1,2,3,4)]) and an empty product is interpreted as 1 and the parameters are such that no pole of $\Gamma(b_j - s), j = 1, 2, 3, \dots, m$ coincides with any pole of $\Gamma(1 - a_k + s), k = 1, 2, 3, \dots, n$. Without any loss of generality, we are assuming that $p \leq q$.

Translation property [7, p.1066, Entry(24)]

$$(1.6) \quad z^\sigma G_{p,q}^{m,n} \left(z \left| \begin{array}{c} a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m, b_{m+1}, \dots, b_q \end{array} \right. \right) = \\ = G_{p,q}^{m,n} \left(z \left| \begin{array}{c} a_1 + \sigma, a_2 + \sigma, \dots, a_n + \sigma, a_{n+1} + \sigma, \dots, a_p + \sigma \\ b_1 + \sigma, b_2 + \sigma, \dots, b_m + \sigma, b_{m+1} + \sigma, \dots, b_q + \sigma \end{array} \right. \right).$$

Symmetric property (Transformation formula)

$$(1.7) \quad G_{p,q}^{m,n} \left(z \left| \begin{array}{c} a_1, a_2, \dots, a_n, a_{n+1}, \dots, a_p \\ b_1, b_2, \dots, b_m, b_{m+1}, \dots, b_q \end{array} \right. \right) = \\ = G_{q,p}^{n,m} \left(\frac{1}{z} \left| \begin{array}{c} 1 - b_1, 1 - b_2, \dots, 1 - b_m, 1 - b_{m+1}, \dots, 1 - b_q \\ 1 - a_1, 1 - a_2, \dots, 1 - a_n, 1 - a_{n+1}, \dots, 1 - a_p \end{array} \right. \right).$$

Reduction formula between ${}_pF_q$ and G-functions [3, p.215, Entry(5.6.1)]; [15, p.47, Entry(9)], is given by when $p \leq q + 1$, then

$$(1.8) \quad \frac{\Gamma(a_1)\Gamma(a_2)\dots\Gamma(a_p)}{\Gamma(b_1)\Gamma(b_2)\dots\Gamma(b_q)} {}_pF_q \left[\begin{array}{c} a_1, a_2, \dots, a_p; \\ b_1, b_2, \dots, b_q; \end{array} z \right] = G_{p,q+1}^{1,p} \left(-z \left| \begin{array}{c} 1 - a_1, 1 - a_2, \dots, 1 - a_p \\ 0, 1 - b_1, 1 - b_2, \dots, 1 - b_q \end{array} \right. \right) \\ = G_{q+1,p}^{p,1} \left(-\frac{1}{z} \left| \begin{array}{c} 1, b_1, b_2, \dots, b_q \\ a_1, a_2, \dots, a_p \end{array} \right. \right), \\ \left(p \leq q \text{ and } |z| < \infty, \text{ or } p = q + 1 \text{ and } |z| < 1 \right).$$

2. MAIN RESULT AND ITS DEMONSTRATION

Our main result (presumably new) is stated here as the following theorem. The result derived in this section is interesting and potentially useful.

The following hypergeometric summation theorem holds true, when any values of parameters and variables leading to the result which do not make sense, are tacitly excluded and each of the series involved in the following result is absolutely convergent.

Then

Theorem 2.1.

$${}_{p+1}F_q \left[\begin{array}{c} a + m, \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{array} z \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(\alpha_1)_r (\alpha_2)_r \dots (\alpha_p)_r z^r}{(\beta_1)_r (\beta_2)_r \dots (\beta_q)_r} \times \right.$$

$$(2.1) \quad \left. \times_{p+1} F_q \left[\begin{matrix} a+r, \alpha_1+r, \alpha_2+r, \dots, \alpha_p+r; \\ \beta_1+r, \beta_2+r, \dots, \beta_q+r; \end{matrix} z \right] \right\},$$

$\left(a, \alpha_i, \beta_j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; \{i=1,2,3,\dots,p ; j=1,2,3,\dots,q\} ; p=q, |z| < 1, \right.$
 $\text{or } p \leq (q-1), |z| < \infty, \text{ or } p=q, z=1, \Re(-a-m + \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i) > 0,$
 $\left. \text{or } p=q, z=-1, \Re(-a-m + \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i) > -1 ; m \in \mathbb{N}_0 \right).$

Demonstration : In order to derive the theorem (2.1) we consider the following Mellin-Barnes type contour integral

$$\begin{aligned} & \times_{p+1} F_q \left[\begin{matrix} a-m, \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{matrix} z \right] = \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a-m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \times \\ & \times \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} \frac{\Gamma(-s)\Gamma(a-m+s)\Gamma(\alpha_1+s)\Gamma(\alpha_2+s)\dots\Gamma(\alpha_p+s)(-z)^s ds}{\Gamma(\beta_1+s)\Gamma(\beta_2+s)\dots\Gamma(\beta_q+s)} \\ & = \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a-2m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \times \\ & \times \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} \frac{\Gamma(-s)\Gamma(a-2m+s)\Gamma(\alpha_1+s)\Gamma(\alpha_2+s)\dots\Gamma(\alpha_p+s)}{\Gamma(\beta_1+s)\Gamma(\beta_2+s)\dots\Gamma(\beta_q+s)} {}_2F_1 \left[\begin{matrix} -m, -s; \\ a-2m; \end{matrix} 1 \right] (-z)^s ds \\ & = \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a-2m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \times \\ & \times \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} \frac{\Gamma(-s)\Gamma(a-2m+s)\Gamma(\alpha_1+s)\Gamma(\alpha_2+s)\dots\Gamma(\alpha_p+s)}{\Gamma(\beta_1+s)\Gamma(\beta_2+s)\dots\Gamma(\beta_q+s)} \sum_{r=0}^m \left\{ \frac{(-m)_r (-s)_r}{(a-2m)_r r!} \right\} (-z)^s ds \\ & = \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a-2m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \sum_{r=0}^m \left\{ \frac{(-m)_r}{(a-2m)_r r!} \right\} \times \\ & \times \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} \frac{\Gamma(r-s)\Gamma(1-(1-a+2m)+s)\Gamma(1-(1-\alpha_1)+s)\dots\Gamma(1-(1-\alpha_p)+s)(-z)^s ds}{\Gamma(1-(1-\beta_1)+s)\Gamma(1-(1-\beta_2)+s)\dots\Gamma(1-(1-\beta_q)+s)} \\ & = \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a-2m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \times \\ & \times \sum_{r=0}^m \left\{ \frac{(-m)_r}{(a-2m)_r r!} G_{p+1,q+1}^{1,p+1} \left(-z \left| \begin{matrix} 1-a+2m, 1-\alpha_1, 1-\alpha_2, \dots, 1-\alpha_p \\ r, 1-\beta_1, 1-\beta_2, \dots, 1-\beta_q \end{matrix} \right. \right) \right\} \end{aligned}$$

Now using the symmetric property (1.7), we get

$$\begin{aligned} {}_{p+1}F_q \left[\begin{matrix} a - m, \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{matrix} z \right] &= \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a - 2m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \times \\ &\times \sum_{r=0}^m \left\{ \frac{(-m)_r (-z)^r}{(a - 2m)_r r!} \left(\frac{-1}{z} \right)^r G_{q+1, p+1}^{p+1, 1} \left(-\frac{1}{z} \middle| \begin{matrix} 1 - r, \beta_1, \beta_2, \dots, \beta_q \\ a - 2m, \alpha_1, \alpha_2, \dots, \alpha_p \end{matrix} \right) \right\} \end{aligned}$$

Now applying the translation property (1.6), we get

$$\begin{aligned} {}_{p+1}F_q \left[\begin{matrix} a - m, \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{matrix} z \right] &= \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a - 2m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \times \\ &\times \sum_{r=0}^m \left\{ \frac{(-m)_r (-z)^r}{(a - 2m)_r r!} G_{q+1, p+1}^{p+1, 1} \left(-\frac{1}{z} \middle| \begin{matrix} 1, \beta_1 + r, \beta_2 + r, \dots, \beta_q + r \\ a - 2m + r, \alpha_1 + r, \alpha_2 + r, \dots, \alpha_p + r \end{matrix} \right) \right\} \end{aligned}$$

Now using the reduction formula (1.8), we get

$$\begin{aligned} {}_{p+1}F_q \left[\begin{matrix} a - m, \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{matrix} z \right] &= \frac{\Gamma(\beta_1)\Gamma(\beta_2)\dots\Gamma(\beta_q)}{\Gamma(a - 2m)\Gamma(\alpha_1)\Gamma(\alpha_2)\dots\Gamma(\alpha_p)} \times \\ &\times \sum_{r=0}^m \left\{ \frac{(-m)_r (-z)^r}{(a - 2m)_r r!} \frac{\Gamma(a - 2m + r)\Gamma(\alpha_1 + r)\Gamma(\alpha_2 + r)\dots\Gamma(\alpha_p + r)}{\Gamma(\beta_1 + r)\Gamma(\beta_2 + r)\dots\Gamma(\beta_q + r)} \times \right. \\ &\quad \left. \times {}_{p+1}F_q \left[\begin{matrix} a - 2m + r, \alpha_1 + r, \alpha_2 + r, \dots, \alpha_p + r; \\ \beta_1 + r, \beta_2 + r, \dots, \beta_q + r; \end{matrix} z \right] \right\} \end{aligned}$$

Therefore

$$\begin{aligned} (2.2) \quad {}_{p+1}F_q \left[\begin{matrix} a - m, \alpha_1, \alpha_2, \dots, \alpha_p; \\ \beta_1, \beta_2, \dots, \beta_q; \end{matrix} z \right] &= \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(\alpha_1)_r (\alpha_2)_r \dots (\alpha_p)_r z^r}{(\beta_1)_r (\beta_2)_r \dots (\beta_q)_r} \times \right. \\ &\quad \left. \times {}_{p+1}F_q \left[\begin{matrix} a - 2m + r, \alpha_1 + r, \alpha_2 + r, \dots, \alpha_p + r; \\ \beta_1 + r, \beta_2 + r, \dots, \beta_q + r; \end{matrix} z \right] \right\} \\ &\left(a - 2m, \alpha_i, \beta_j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; \{i=1, 2, 3, \dots, p ; j=1, 2, 3, \dots, q\} ; p = q, |z| < 1, \right. \\ &\text{or } p \leq (q - 1), |z| < \infty, \text{ or } p = q, z = 1, \Re(-a + m + \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i) > 0, \\ &\left. \text{or } p = q, z = -1, \Re(-a + m + \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i) > -1 ; m \in \mathbb{N}_0 \right). \end{aligned}$$

Replace a by $a + 2m$ in equation (2.2), we get the theorem (2.1).

3. APPLICATIONS IN CLAUSEN SERIES

In this section, we obtain some special cases of our main summation theorem (2.1)

In theorem (2.1) put $p = 2$ and $q = 2$, we get

$$(3.1) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ g, h; \end{matrix} z \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r (c)_r (z)^r}{(g)_r (h)_r} {}_3F_2 \left[\begin{matrix} a+r, b+r, c+r; \\ g+r, h+r; \end{matrix} z \right] \right\},$$

$$\left(a, b, c, g, h \in \mathbb{C} \setminus \mathbb{Z}_0^-; |z| < 1; m \in \mathbb{N}_0 \right).$$

In equation (3.1) put $g = a$ and $z = -1$, we get

$$(3.2) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, h; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r (c)_r (-1)^r}{(a)_r (h)_r} {}_2F_1 \left[\begin{matrix} b+r, c+r; \\ h+r; \end{matrix} -1 \right] \right\},$$

$$\left(\Re(h-b-c) > m-1; a, b, c, h \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).$$

In equation (3.2) put $h = 1+b-c-p$ and use the summation theorem [2, p.1523, Equation (2.2)] of Choi-Rathie and Malani, we get

$$(3.3) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, 1+b-c-p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(c)_r \Gamma(1+b-c-p)}{2(-1)^r \Gamma(b)(a)_r} \sum_{k=0}^{p-r} \left[\binom{p-r}{k} \frac{\Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{b+r+k+2-2c-2p}{2})} \right] \right\},$$

$$\left(\Re(c) < \frac{2-p-m}{2}; a, b, c, 1+b-c-p \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0; p \geq m \right).$$

In equation (3.2) put $h = 1+b-c+p$ and use the summation theorem [2, p.1524, Equation (2.3)] of Choi-Rathie and Malani, we get

$$(3.4) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, 1+b-c+p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{\Gamma(1+b-c+p)}{2\Gamma(b)(a)_r(1-c)_p} \sum_{k=0}^{p+r} \left[\binom{p+r}{k} \frac{(-1)^k \Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{b-r+k+2-2c}{2})} \right] \right\},$$

$$\left(\Re(c) < \frac{p-m+2}{2}; a, b, c, 1-c-m, 1+b-c \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0 \right).$$

In equation (3.2) put $h = b-c-p$ and use the summation theorem [10, p.14, Equation (3.1)] of authors, we get

$${}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, b-c-p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{\Gamma(b-c-p)(c)_r (-1)^r}{2(a)_r \Gamma(b)} \times \right.$$

$$(3.5) \quad \times \sum_{k=0}^{p-r} \binom{p-r}{k} \left[\frac{\Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{b+r+k-2c-2p}{2})} + \frac{\Gamma(\frac{b+r+k+1}{2})}{\Gamma(\frac{b+r+k+1-2c-2p}{2})} \right] \Bigg\} ,$$

$$\left(\Re(c) < \frac{1-m-p}{2} ; a, b, c, b-c-p \in \mathbb{C} \setminus \mathbb{Z}_0^- ; m, p \in \mathbb{N}_0 ; p \geq m \right).$$

In equation (3.2) put $h = b - c + p$ and use the summation theorem [10, p.14, Equation (3.2)] of authors, we get

$$(3.6) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, b-c+p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{\Gamma(b-c+p)(c)_r}{2(a)_r(c+1)_r(-c)_p \Gamma(b)} \times \right.$$

$$\times \sum_{k=0}^{p+r} \binom{p+r}{k} \left[\frac{(-1)^k \Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{b-r+k-2c}{2})} + \frac{(-1)^k \Gamma(\frac{b+r+k+1}{2})}{\Gamma(\frac{b-r+k+1-2c}{2})} \right] \Bigg\} ,$$

$$\left(\Re(c) < \frac{p+1-m}{2} ; a, b, c, -c-m, b-c \in \mathbb{C} \setminus \mathbb{Z}_0^- ; m, p \in \mathbb{N}_0 \right).$$

When $g = a$ and $z = \frac{1}{2}$ in equation (3.1), we get

$$(3.7) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, h; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r(c)_r}{2^r (a)_r(h)_r} {}_2F_1 \left[\begin{matrix} b+r, c+r; \\ h+r; \end{matrix} \frac{1}{2} \right] \right\} ,$$

$$\left(a, b, c, h \in \mathbb{C} \setminus \mathbb{Z}_0^- ; m \in \mathbb{N}_0 \right).$$

In equation (3.7) put $h = \frac{b+c+1-p}{2}$ and use the summation theorem [9, p.491, Entry (7.3.7.2)] of Prudnikov *et al.*, we get

$$(3.8) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, \frac{b+c+1-p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{2^{c-1} (b)_r \Gamma(\frac{b+c+1-p}{2})}{(a)_r \Gamma(c)} \sum_{k=0}^p \left[\binom{p}{k} \frac{\Gamma(\frac{c+r+k}{2})}{\Gamma(\frac{b+r+k-p+1}{2})} \right] \right\} ,$$

$$\left(a, b, c, \frac{b+c+1-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^- ; m, p \in \mathbb{N}_0 \right).$$

In equation (3.7) put $h = \frac{b+c+1+p}{2}$ and use the summation theorem [14, p.827, Theorem 1] of Rakha-Rathie, we get

$$(3.9) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, \frac{b+c+1+p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{2^{c-1} (b)_r \Gamma(\frac{b+c+1+p}{2})}{(a)_r \Gamma(c) (\frac{b-c+1-p}{2})_p} \sum_{k=0}^p \left[\binom{p}{k} \frac{(-1)^k \Gamma(\frac{c+r+k}{2})}{\Gamma(\frac{b+r+k-p+1}{2})} \right] \right\} ,$$

$$\left(a, b, c, \frac{1+b+c+p}{2}, \frac{b-c+1-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^- ; p, m \in \mathbb{N}_0 \right).$$

In equation (3.7) put $h = \frac{b+c-p}{2}$ and use the summation theorem [11, p.48, Equation (3.1)] of authors, we get

$$(3.10) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, \frac{b+c-p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{2^{b-1}(c)_r \Gamma(\frac{b+c-p}{2})}{(a)_r \Gamma(b)} \sum_{k=0}^p \binom{p}{k} \left[\frac{\Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{c+r+k-p}{2})} + \frac{\Gamma(\frac{b+r+k+1}{2})}{\Gamma(\frac{c+r+k-p+1}{2})} \right] \right\},$$

$$\left(a, b, c, \frac{b+c-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; p, m \in \mathbb{N}_0 \right).$$

In equation (3.7) put $h = \frac{b+c+p}{2}$ and use the summation theorem [1, p.582, Entry (8.1.1.130)] of Brychkov, we get

$$(3.11) \quad {}_3F_2 \left[\begin{matrix} a+m, b, c; \\ a, \frac{b+c+p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r (c)_r \Gamma(\frac{1}{2}) \Gamma(\frac{b+c+p}{2})}{2^r (a)_r (\frac{-b+c-p}{2})_p} \times \right.$$

$$\left. \times \sum_{k=0}^p \left(\binom{p}{k} (-2)^{-k} (b+r)_k \left[\frac{1}{\Gamma(\frac{b+r+k+1}{2}) \Gamma(\frac{c+r+k-p}{2})} + \frac{1}{\Gamma(\frac{b+r+k}{2}) \Gamma(\frac{c+r+k-p+1}{2})} \right] \right) \right\},$$

$$\left(a, b, c, \frac{b+c+p}{2}, \frac{-b+c-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; p, m \in \mathbb{N}_0 \right).$$

In equation (3.7) put $c = 1 - b - p$ and use the summation theorem [14, p.828, Theorem (6)] of Rakha-Rathie, we get

$$(3.12) \quad {}_3F_2 \left[\begin{matrix} a+m, b, 1-b-p; \\ a, h; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r (1-b-p)_r \Gamma(h)}{(a)_r \Gamma(h-b) 2^{b+p}} \times \right.$$

$$\left. \times \sum_{k=0}^{p-2r} \left[\binom{p-2r}{k} \frac{\Gamma(\frac{h-b+k}{2})}{\Gamma(\frac{h+b+k+2r}{2})} \right] \right\},$$

$$\left(a, b, 1-b-p, h, h-b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0; p \geq 2m \right).$$

In equation (3.7) put $c = 1 - b + p$ and use the summation theorem [14, p.827, Theorem (5)] of Rakha-Rathie, we get

$$(3.13) \quad {}_3F_2 \left[\begin{matrix} a+m, b, 1-b+p; \\ a, h; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(1-b+p)_r \Gamma(h) \Gamma(b-r-p)}{(a)_r \Gamma(b) \Gamma(h-b) 2^{b-p}} \times \right.$$

$$\left. \times \sum_{k=0}^{p+2r} \left[\binom{p+2r}{k} \frac{(-1)^k \Gamma(\frac{h-b+k}{2})}{\Gamma(\frac{h+b+k-2r-2p}{2})} \right] \right\},$$

$$\left(a, b, 1-b, h, h-b, b-m-p \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0 \right).$$

In equation (3.7) put $c = -b - p$ and use the summation theorem [12, Equation (3.3)] of authors, we get

$$(3.14) \quad {}_3F_2 \left[\begin{matrix} a + m, b, -b - p; \\ a, h; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r (-b-p)_r \Gamma(h) 2^{-b-p-1}}{(a)_r \Gamma(h-b)} \times \right. \\ \left. \times \sum_{k=0}^{p-2r} \binom{p-2r}{k} \left[\frac{\Gamma(\frac{h-b+k}{2})}{\Gamma(\frac{h+b+k+2r}{2})} + \frac{\Gamma(\frac{h-b+k+1}{2})}{\Gamma(\frac{h+b+k+2r+1}{2})} \right] \right\}, \\ \left(a, b, -b-p, h, h-b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0; p \geq 2m \right).$$

In equation (3.7) put $c = -b + p$ and use the summation theorem [12, Equation (3.5)] of authors, we get

$$(3.15) \quad {}_3F_2 \left[\begin{matrix} a + m, b, -b + p; \\ a, h; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(-b+p)_r \Gamma(h) \Gamma(b-p-r) 2^{p-b-1}}{(a)_r \Gamma(b) \Gamma(h-b)} \times \right. \\ \left. \times \sum_{k=0}^{p+2r} \binom{p+2r}{k} \left[\frac{(-1)^k \Gamma(\frac{h-b+k}{2})}{\Gamma(\frac{h+b+k-2p-2r}{2})} + \frac{(-1)^k \Gamma(\frac{h-b+k+1}{2})}{\Gamma(\frac{h+b+k-2r-2p+1}{2})} \right] \right\}, \\ \left(a, b, -b, b-m-p, h, h-b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0 \right).$$

In equation (3.1) put $g = a$ and $z = 1$ and use the Classical Gauss summation theorem [15, p.30, Equation 1.2(7)], we get

$$(3.16) \quad {}_3F_2 \left[\begin{matrix} a + m, b, c; \\ a, h; \end{matrix} 1 \right] = \frac{\Gamma(h) \Gamma(h-b-c)}{\Gamma(h-b) \Gamma(h-c)} {}_3F_2 \left[\begin{matrix} -m, b, c; \\ a, 1+b+c-h; \end{matrix} 1 \right], \\ \left((h-b-c) \neq 0, \pm 1, \pm 2, \dots; \Re(h-b-c) > m; \right. \\ \left. a, b, c, h, 1+b+c-h \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).$$

Now put $h = c + 1$ in the equation (3.16), we get

$$(3.17) \quad {}_3F_2 \left[\begin{matrix} a + m, b, c; \\ a, c + 1; \end{matrix} 1 \right] = \frac{\Gamma(c+1) \Gamma(1-b) (a-c)_m}{\Gamma(c+1-b) (a)_m}, \\ \left((1-b) \neq 0, \pm 1, \pm 2, \dots; \Re(1-b) > m; a, b, c \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).$$

which is a known summation theorem of Mitra [8, p.85, Last Equation].

4. APPLICATIONS IN GAUSSIAN SERIES

In equation (3.1) put $h = c$ and $z = -1$, we get

$$(4.1) \quad {}_2F_1 \left[\begin{matrix} a+m, b; \\ g; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r (-1)^r}{(g)_r} {}_2F_1 \left[\begin{matrix} a+r, b+r; \\ g+r; \end{matrix} -1 \right] \right\},$$

$$\left(\Re(g-a-b) > m-1; a, b, g \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).$$

In equation (4.1) put $g = 1+a-b-p$ and use the summation theorem [2, p.1523, Equation (2.2)] of Choi-Rathie and Malani, we get

$$(4.2) \quad {}_2F_1 \left[\begin{matrix} a+m, b; \\ 1+a-b-p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r \Gamma(1+a-b-p)}{2(-1)^r \Gamma(a+r)} \sum_{k=0}^{p-r} \left[\binom{p-r}{k} \frac{\Gamma(\frac{a+r+k}{2})}{\Gamma(\frac{a+r+k+2-2b-2p}{2})} \right] \right\},$$

$$\left(\Re(b) < \frac{2-p-m}{2}; a, b, 1+a-b-p \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0; p \geq m \right).$$

In equation (4.1) put $g = 1+a-b+p$ and use the summation theorem [2, p.1524, Equation (2.3)] of Choi-Rathie and Malani, we get

$$(4.3) \quad {}_2F_1 \left[\begin{matrix} a+m, b; \\ 1+a-b+p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{\Gamma(1+a-b+p)}{2(1-b)_p \Gamma(a+r)} \sum_{k=0}^{p+r} \left[\binom{p+r}{k} \frac{(-1)^k \Gamma(\frac{a+r+k}{2})}{\Gamma(\frac{a-r+k+2-2b}{2})} \right] \right\},$$

$$\left(\Re(b) < \frac{p+2-m}{2}; a, b, 1-b-m, 1+a-b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0 \right).$$

In equation (4.1) put $g = a-b-p$ and use the summation theorem [10, p.14, Equation (3.1)] of authors, we get

$$(4.4) \quad {}_2F_1 \left[\begin{matrix} a+m, b; \\ a-b-p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r \Gamma(a-b-p) (-1)^r}{2\Gamma(a+r)} \times \right.$$

$$\left. \times \sum_{k=0}^{p-r} \binom{p-r}{k} \left[\frac{\Gamma(\frac{a+r+k}{2})}{\Gamma(\frac{a+r+k-2b-2p}{2})} + \frac{\Gamma(\frac{a+r+k+1}{2})}{\Gamma(\frac{a+r+k+1-2b-2p}{2})} \right] \right\},$$

$$\left(\Re(b) < \frac{1-p-m}{2}; a, b, a-b-p \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0; p \geq m \right).$$

In equation (4.1) put $g = a-b+p$ and use the summation theorem [10, p.14, Equation (3.2)] of authors, we get

$$\begin{aligned}
& {}_2F_1 \left[\begin{matrix} a+m, b; \\ a-b+p; \end{matrix} -1 \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r \Gamma(a-b+p)}{2(-b)_p (1+b)_r \Gamma(a+r)} \times \right. \\
(4.5) \quad & \left. \times \sum_{k=0}^{p+r} \binom{p+r}{k} \left[\frac{(-1)^k \Gamma(\frac{a+r+k}{2})}{\Gamma(\frac{a-r+k-2b}{2})} + \frac{(-1)^k \Gamma(\frac{a+r+k+1}{2})}{\Gamma(\frac{a-r+k+1-2b}{2})} \right] \right\}, \\
& \left(\Re(b) < \frac{p+1-m}{2}; a, b, -b-m, a-b \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0 \right).
\end{aligned}$$

In equation (3.1) put $h = c$ and $z = \frac{1}{2}$, we get

$$\begin{aligned}
(4.6) \quad & {}_2F_1 \left[\begin{matrix} a+m, b; \\ g; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r}{2^r (g)_r} {}_2F_1 \left[\begin{matrix} a+r, b+r; \\ g+r; \end{matrix} \frac{1}{2} \right] \right\}, \\
& \left(a, b, g \in \mathbb{C} \setminus \mathbb{Z}_0^-; m \in \mathbb{N}_0 \right).
\end{aligned}$$

In equation (4.6) put $g = \frac{a+b+1-p}{2}$ and use the summation theorem [9, p.491, Entry (7.3.7.2)] of Prudnikov *et al.*, we get

$$\begin{aligned}
(4.7) \quad & {}_2F_1 \left[\begin{matrix} a+m, b; \\ \frac{a+b+1-p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{2^{b-1} \Gamma(\frac{a+b+1-p}{2})}{\Gamma(b)} \sum_{k=0}^p \left[\binom{p}{k} \frac{\Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{a+r+k+1-p}{2})} \right] \right\}, \\
& \left(a, b, \frac{1+a+b-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; p, m \in \mathbb{N}_0 \right).
\end{aligned}$$

In equation (4.6) put $g = \frac{a+b+1+p}{2}$ and use the summation theorem [14, p.827, Theorem (1)] of Rakha-Rathie, we get

$$\begin{aligned}
(4.8) \quad & {}_2F_1 \left[\begin{matrix} a+m, b; \\ \frac{a+b+1+p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{2^{b-1} \Gamma(\frac{a+b+1+p}{2})}{\Gamma(b) (\frac{a-b+1-p}{2})_p} \sum_{k=0}^p \left[\binom{p}{k} \frac{(-1)^k \Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{a+r+k+1-p}{2})} \right] \right\}, \\
& \left(a, b, \frac{1+a+b+p}{2}, \frac{a-b+1-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; p, m \in \mathbb{N}_0 \right).
\end{aligned}$$

In equation (4.6) put $g = \frac{a+b-p}{2}$ and use the summation theorem [11, p.48, Equation (16)] of authors, we get

$$\begin{aligned}
(4.9) \quad & {}_2F_1 \left[\begin{matrix} a+m, b; \\ \frac{a+b-p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{2^{b-1} \Gamma(\frac{a+b-p}{2})}{\Gamma(b)} \sum_{k=0}^p \binom{p}{k} \left[\frac{\Gamma(\frac{b+r+k}{2})}{\Gamma(\frac{a+r+k-p}{2})} + \frac{\Gamma(\frac{b+r+k+1}{2})}{\Gamma(\frac{a+r+k+1-p}{2})} \right] \right\}, \\
& \left(a, b, \frac{a+b-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; p, m \in \mathbb{N}_0 \right).
\end{aligned}$$

In equation (4.6) put $g = \frac{a+b+p}{2}$ and use the summation theorem [1, p.582, Entry (8.1.1.130)] of Brychkov, we get

$$(4.10) \quad {}_2F_1 \left[\begin{matrix} a+m, b; \\ \frac{a+b+p}{2}; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(b)_r \Gamma(\frac{a+b+p}{2}) \Gamma(\frac{1}{2})}{2^r (\frac{b-a-p}{2})_p} \sum_{k=0}^p \left(\binom{p}{k} (-2)^{-k} (a+r)_k \times \right. \right. \\ \left. \left. \times \left[\frac{1}{\Gamma(\frac{a+r+k+1}{2}) \Gamma(\frac{b+r+k-p}{2})} + \frac{1}{\Gamma(\frac{a+r+k}{2}) \Gamma(\frac{b+r+k+1-p}{2})} \right] \right) \right\}, \\ \left(a, b, \frac{a+b+p}{2}, \frac{-a+b-p}{2} \in \mathbb{C} \setminus \mathbb{Z}_0^-; p, m \in \mathbb{N}_0 \right).$$

In equation (4.6) put $b = 1 - a - p$ and use the summation theorem [14, p.828, Theorem (6)] of Rakha-Rathie, we get

$$(4.11) \quad {}_2F_1 \left[\begin{matrix} a+m, 1-a-p; \\ g; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(1-a-p)_r \Gamma(g)}{\Gamma(g-a) 2^{a+p}} \times \right. \\ \left. \times \sum_{k=0}^{p-2r} \left[\binom{p-2r}{k} \frac{\Gamma(\frac{g-a+k}{2})}{\Gamma(\frac{g+a+k+2r}{2})} \right] \right\}, \\ \left(a, 1-a-p, g, g-a \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0; p \geq 2m \right).$$

In equation (4.6) put $b = 1 - a + p$ and use the summation theorem [14, p.827, Theorem (5)] of Rakha-Rathie, we get

$$(4.12) \quad {}_2F_1 \left[\begin{matrix} a+m, 1-a+p; \\ g; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(1-a+p)_r \Gamma(g) \Gamma(a-r-p)}{\Gamma(a+r) \Gamma(g-a) 2^{a-p}} \times \right. \\ \left. \times \sum_{k=0}^{p+2r} \left[\binom{p+2r}{k} \frac{(-1)^k \Gamma(\frac{g-a+k}{2})}{\Gamma(\frac{g+a+k-2r-2p}{2})} \right] \right\}, \\ \left(a, 1-a, g, g-a, a-m-p \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0 \right).$$

In equation (4.6) put $b = -a - p$ and use the summation theorem [12, Equation (3.3)] of authors, we get

$$(4.13) \quad {}_2F_1 \left[\begin{matrix} a+m, -a-p; \\ g; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(-a-p)_r \Gamma(g) 2^{-a-p-1}}{\Gamma(g-a)} \times \right. \\ \left. \times \sum_{k=0}^{p-2r} \binom{p-2r}{k} \left[\frac{\Gamma(\frac{g-a+k}{2})}{\Gamma(\frac{g+a+k+2r}{2})} + \frac{\Gamma(\frac{g-a+k+1}{2})}{\Gamma(\frac{g+a+k+2r+1}{2})} \right] \right\}, \\ \left(a, -a-p, g, g-a \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0; p \geq 2m \right).$$

In equation (4.6) put $b = -a + p$ and use the summation theorem [12, Equation (3.5)] of authors, we get

$$(4.14) \quad {}_2F_1 \left[\begin{matrix} a + m, -a + p; \\ g; \end{matrix} \frac{1}{2} \right] = \sum_{r=0}^m \left\{ \binom{m}{r} \frac{(-a+p)_r \Gamma(g) \Gamma(a-p-r) 2^{p-a-1}}{\Gamma(a+r) \Gamma(g-a)} \times \right. \\ \left. \times \sum_{k=0}^{p+2r} \binom{p+2r}{k} \left[\frac{(-1)^k \Gamma(\frac{g-a+k}{2})}{\Gamma(\frac{g+a+k-2p-2r}{2})} + \frac{(-1)^k \Gamma(\frac{g-a+k+1}{2})}{\Gamma(\frac{g+a+k-2r-2p+1}{2})} \right] \right\}, \\ \left(a, -a, a-p-m, g, g-a \in \mathbb{C} \setminus \mathbb{Z}_0^-; m, p \in \mathbb{N}_0 \right).$$

We conclude our present investigation by observing that several other corollaries and consequences of hypergeometric summation theorems (2.1), (3.1), (3.2), (3.7), (4.1) and (4.6) can also be deduced in an analogous manner.

REFERENCES

- [1] Brychkov, Yury A. ; *Handbook of Special Functions : Derivatives, Integrals, Series and Other Formulas*, A Chapman & Hall book, CRC Press (Taylor and Francis Group), Boca Raton, London, New York, 2008.
- [2] Choi, J., Rathie, A. K. and Malani, S. ; Kummer's Theorem and its Contiguous Identities, *Taiwanese Journal of Mathematics* 11 (5) (2007), 1521–1527.
- [3] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G. ; *Higher Transcendental Functions*, Vol. I, McGraw -Hill Book Company, New York, Toronto and London, 1953.
- [4] Luke, Y. L. ; *The Special Functions and Their Approximations*, Vol.I, Academic Press, London, 1969.
- [5] Mathai, A. M. and Saxena, R. K.; *Generalized Hypergeometric Functions with Applications in Statistics and Physical Sciences, Lectures Notes in Mathematics No. 348*, Springer-Verlag, Berlin, Heidelberg, New York, 1973.
- [6] Meijer, C. S.; Über Whittakersche Bezw. Besselsche Funktionen und deren Produkte, *Nieuw. Arch. Wiskunde*, 18(2)(1936), 10-39.
- [7] Meijer, C. S.; Multiplikationstheoreme für die Funktion $G_{p,q}^{m,n}(z)$, *Nederl. Akad. Wetensch. Proc. Ser. A*, 44(1941), 1062-1070.
- [8] Mitra, S. C. ; On a Certain Transformation About Generalized Hypergeometric Series, *J. Indian Math. Soc. (N.S.)*, 6(1942), 84–86.
- [9] Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O. I. ; *Integrals and Series, Volume III : More Special Functions*, Nauka, Moscow, 1986 (In Russian); (Translated from the Russian by G.G.Gould) Gordon and Breach Science Publishers, New York, 1990.
- [10] Qureshi, M. I. and Baboo, M. S. ; Some Unified and Generalized Kummer's First Summation Theorems with Applications in Laplace Transform Technique, *Asia Pacific Journal of Mathematics*. 3 (1) (2016), 10–23.
- [11] Qureshi, M. I. and Baboo, M. S. ; Some Unified and Generalized Kummer's Second Summation Theorems with Applications in Laplace Transform Technique, *International Journal of Mathematics and its Applications* 4(1-C), (2016) 45–52.

- [12] Qureshi, M. I. and Baboo, M. S. ; Some Unified and Generalized Kummer's Third Summation Theorems with Applications in Laplace Transform Technique, First International Conference cum Exhibition on Building Utility, Faculty of Engineering and Technology Jamia Millia Islamia, New Delhi, India, ICEBU, 1-3 December- 2016, pp.139–150. Enriched Publications Pvt.Ltd. New Delhi 110075 India.
- [13] Rainville, E. D. ; *Special Functions*, The Macmillan Company, New York, 1960 ; Reprinted by Chelsea Publ. Co., Bronx, New York, 1971.
- [14] Rakha, M. A. and Rathie, A. K. ; Generalizations of Classical Summation Theorems for the Series ${}_2F_1$ and ${}_3F_2$ with Applications, *Integral Transforms and Special Functions*, 22 (11) (2011), 823–840.
- [15] Srivastava, H. M. and Manocha, H. L. ; *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1984.